BULLETIN OF THE AMERICAN MATHEMATICAL SOCIETY.

CONTINUATION OF THE BULLETIN OF THE NEW YORK MATHEMATICAL SOCIETY.

A HISTORICAL AND CRITICAL REVIEW OF MATHEMATICAL SCIENCE

EDITED BY

F. N. COLE, ALEXANDER ZIWET, F. MORLEY,
E. O. LOVETT, C. L. BOUTON,
D. E. SMITH.

VOL. VIII.

OCTOBER 1901 TO JULY 1902.

PUBLISHED FOR THE SOCIETY.
BY
THE MACMILLAN COMPANY,
LANCASTER, PA., AND NEW YORK,
1902.

Reprinted 1968

2ND SERIES, VOL. VIII. No. 1

OCTOBER, 1901

Whole No. 101

CONTENTS

PAGE

The Eighth Summer Meeting of the American Mathematical Society. By Professor F. N. COLE, 1

The Ithaca Colloquium. By Dr. EDWARD KASNER, 22

Upon the Non-Isomorphism of two Simple Groups of Order 8!/2. By Miss IDA M. SCHOTTENFELS, 25

Concerning Surfaces whose First and Second Fundamental Forms are the Second and First Fundamental Forms Respectively of Another Surface. By Professor ALEXANDER PELL, 27

Notes, 29

New Publications, 39

2ND SERIES, VOL. VIII. No. 2

NOVEMBER, 1901

Whole No. 102

CONTENTS

PAGE

On Wronskians of Functions of a Real Variable. By Professor MAXIME BÔCHER, 53

The Configurations of the 27 Lines on a Cubic Surface and the 28 Bitangents to a Quartic Curve. By Professor L. E. DICKSON, 63

The Fiftieth Annual Meeting of the American Association for the Advancement of Science. By Professor G. A. MILLER, 71

Riemann-Weber: Partial Differential Equations of Mathematical Physics. By Professor J. S. AMES 81

Notes, 85

New Publications, 89

2ND SERIES, VOL. VIII. No. 3

DECEMBER, 1901

Whole No. 103

CONTENTS

PAGE

The October Meeting of the American Mathematical Society. By Professor EDWARD KASNER, 95

Modern Methods of Treating Dynamical Problems and in Particular the Problems of Three Bodies. By Professor E. W. BROWN, 103

The Hamburg Meeting of the Deutsche Mathematiker-Vereinigung, September, 1901. By Professor C. M. MASON, 113

Some Curious Properties of Conics Touching the Line Infinity at one of the Circular Points. By Professor E. V. HUNTINGTON and Professor J. K. WHITTEMORE, 122

Picard's Traité d'Analyse. By Professor MAXIME BÔCHER, 124

Errata, 129

Notes, 129

New Publications, 133

2ND SERIES, VOL. VIII. No. 4

JANUARY, 1902

Whole No. 104

CONTENTS

PAGE

Note on Mr. George Peirce's Approximate Construction for π. By Professor M. EMILE LEMOINE, 137

Concerning the Elliptic $\wp(g_2, g_3, z)$-Functions as Coördinates in a Line Complex, and Certain Related Theorems. By Dr. H. F. STECKER, 148

On the Abelian Groups which are Conformal with Non-Abelian Groups. By Professor G. A. MILLER, 154

The Infinitesimal Generators of Certain Parameter Groups. By Dr. S. E. SLOCUM, 156

Shorter Notices, 168

Notes, 172

New Publications, 176

2ND SERIES, VOL. VIII. No. 5

FEBRUARY, 1902

Whole No. 105

CONTENTS

PAGE

The Eighth Annual Meeting of the American Mathematical Society. By Professor F. N. COLE, 183
The January Meeting of the Chicago Section. By Professor THOMAS F. HOLGATE, 198
Vector Analysis. By Professor ALEXANDER ZIWET, 207
Euclid's Elements. By Professor JULIAN LOWELL COOLIDGE, 216
Notes, 221
New Publications, 224

2ND SERIES, VOL. VIII. No. 6

MARCH, 1902

Whole No. 106

CONTENTS

PAGE

The Application of the Fundamental Laws of Algebra to the Multiplication of Infinite Series. By Professor FLORIAN CAJORI, 231
Concerning the Class of a Group of Order p^m that Contains an Operator of Order p^{m-2} or p^{m-3}, p being a Prime. By Dr. W. B. FITE, 236
Proof that the Group of an Irreducible Linear Differential Equation is Transitive. By Dr. SAUL EPSTEEN, 239
Lines of Length Zero on Surfaces. By Dr. L. P. EISENHART, 241
Some Properties of Potential Surfaces. By Dr. EDWARD KASNER, 243
A Modern English Calculus. By Professor W. F. OSGOOD, 248
Shorter Notices, 257
Notes, 261
New Publications, 265

2ND SERIES, VOL. VIII. No. 7

APRIL, 1902

Whole No. 107

CONTENTS

	PAGE
The February Meeting of the American Mathematical Society. By Dr. EDWARD KASNER,	271
Note on the Transformation of a Group into its Canonical Form. By Dr. S. E. SLOCUM,	280
Some Applications of Green's Theorem in One Dimension. By Mr. OTTO DUNKEL,	288
On the Forms of Quintic Scrolls. By Dr. VIRGIL SNYDER,	293
Simplified Definition of a Group. By Dr. E. V. HUNTINGTON,	296
Note on Isotropic Congruences. By Dr. L. P. EISENHART,	301
Kronecker's Lectures on the Theory of Numbers. By Professor G. A. MILLER,	303
Errata,	308
Notes,	308
New Publications,	315

2ND SERIES, VOL. VIII. No. 8

MAY, 1902

Whole No. 108

CONTENTS

	PAGE
The March Meeting of the Chicago Section. By Professor T. F. HOLGATE,	319
Concerning the Angles and the Angular Determination of Planes in 4-Space. By Dr. C. J. KEYSER,	324
Note on the Sufficient Conditions for an Analytic Function. By Mr. D. R. CURTISS,	329
Scheffer's Theory of Surfaces. By Professor J. M. PAGE,	332
Some Recent Books on Mechanics. By Dr. E. B. WILSON,	341
The Galois Theory in Burnside and Panton's Theory of Equations. By Dr. B. S. EASTON,	349
Shorter Notices,	352
Notes,	355
New Publications,	361

2ND SERIES, VOL. VIII. No. 9

JUNE, 1902

Whole No. 109

CONTENTS

	PAGE
The April Meeting of the American Mathematical Society. By Professor F. N. COLE,	367
The Infinitesimal Generators of Parameter Groups. By Professor T. J. I'A. BROMWICH,	375
On the Parabolas (or Paraboloids) Through the Points Common to Two Given Conics (or Quadrics). By Professor T. J. I'A. BROMWICH,	386
A Second Definition of a Group. By Dr. E. V. HUNTINGTON,	388
Determination of All the Groups of Order p^m, p being Any Prime, which Contain the Abelian Group of Order p^{m-1} and of Type $(1, 1, 1, \cdots)$. By Professor G. A. MILLER,	391
A Class of Simply Transitive Linear Groups. By Professor L. E. DICKSON,	394
Errors in Legendre's Tables of Linear Divisors. By Dr. D. N. LEHMER,	401
Some Recent Books on Mechanics. By Dr. E. B. WILSON,	403
Kiepert's Calculus. By Professor E. W. DAVIS,	412
Notes,	419
New Publications,	422

2ND SERIES, VOL. VIII. No. 10

JULY, 1902

Whole No. 110

CONTENTS

	PAGE
The First Meeting of the San Francisco Section of the American Mathematical Society. By Professor E. J. WILCZYNSKI,	429
Mathematical Problems. By Professor DAVID HILBERT,	437
Reply to Mr. J. L. Coolidge's Review of Hill's Euclid. By Professor M. J. M. HILL,	479
Notes,	481
New Publications,	485
Eleventh Annual List of Papers,	491
Index,	497

BULLETIN OF THE

AMERICAN MATHEMATICAL SOCIETY

THE EIGHTH SUMMER MEETING OF THE AMERICAN MATHEMATICAL SOCIETY.

THE Eighth Summer Meeting, supplemented by the Third Colloquium, of the AMERICAN MATHEMATICAL SOCIETY was held at Cornell University, Ithaca, N. Y., through the week August 19–24, 1901. The four sessions of the Summer Meeting proper occupied the first two days of the week. Only an unduly severe compression made it possible to complete the lengthy programme in this period. More time, especially for the discussion of papers, is needed at the larger meetings of the year, and this question is to receive immediate attention. On the present occasion various circumstances made an adequate provision of time practically impossible.

The attendance at the Summer Meeting numbered about sixty persons, including the following forty-five members of the Society:

Dr. G. A. Bliss, Professor Oskar Bolza, Professor E. W. Brown, Dr. W. G. Bullard, Professor F. N. Cole, Professor L. L. Conant, Mr. A. T. DeLury, Professor L. W. Dowling, Professor L. E. Dickson, Professor W. P. Durfee, Professor H. T. Eddy, Professor John Eiesland, Professor Peter Field, Dr. J. C. Fields, Professor T. S. Fiske, Mr. W. B. Ford, Mr. B. F. Groat, Dr. E. R. Hedrick, Professor T. F. Holgate, Dr. J. I. Hutchinson, Dr. Edward Kasner, Mr. C. J. Keyser, Dr. G. H. Ling, Professor T. E. McKinney, Professor James McMahon, Professor W. H. Maltbie, Professor J. L. Markley, Professor W. H. Metzler, Dr. G. A. Miller, Professor E. H. Moore, Dr. G. A. Murray, Professor G. D. Olds, Professor W. F. Osgood, Mr. F. G. Radelfinger, Professor E. D. Roe, Professor E. B. Skinner, Dr. Virgil Snyder, Dr. H. F. Stecker, Professor J. H. Tanner, Professor E. J. Townsend, Professor E. B. Van Vleck, Professor L. A. Wait, Professor L. G. Weld, Dr. J. v. E. Westfall, Professor Alexander Ziwet.

The President of the Society, Professor Eliakim Hastings Moore, occupied the chair, being relieved by Vice-President Professor T. S. Fiske. An address of welcome by Professor L. A. Wait in behalf of the University, was the forerunner of a most generous hospitality accorded the Society by the University and its individual officers. Resolutions adopted at the closing session partially express the Society's sense of appreciation of the courtesies extended to it throughout its stay in Ithaca.

The Council announced the election of the following persons to membership in the Society: Dr. E. R. Hedrick, Yale University; Professor S. W. Reaves, Clemson College, S. C., Twelve applications for admission to the Society were received. A committee consisting of the President, Professor Fiske, and Professor Osgood, was appointed to prepare and submit to the Council at the October meeting a list of nominations of officers of the Society for the coming year.

The following papers were presented at the meeting:

(1) Professor MAXIME BÔCHER: "On certain pairs of transcendental functions whose roots separate each other."

(2) Dr. J. I. HUTCHINSON: "On a class of automorphic functions."

(3) Professor A. PRINGSHEIM: "Ueber den Goursat'schen Beweis des Cauchy'schen Integralsatzes."

(4) Professor A. PRINGSHEIM: "Ueber die Anwendung der Cauchy'schen Multiplicationsregel auf bedingt convergente oder divergente Reihen."

(5) Mr. W. B. FORD: "On the expression of Bessel-functions in terms of the trigonometric functions."

(6) Professor E. H. MOORE: "On the theory of improper definite integrals."

(7) Professor OSKAR BOLZA: "New proof of a theorem of Osgood in the calculus of variations."

(8) Dr. G. A. BLISS: "The problem of the calculus of variations when the end point is variable."

(9) Dr. J. C. FIELDS: "On certain relations existing between the branch points and the double points of an algebraic curve."

(10) Dr. J. C. FIELDS: "The Riemann-Roch theorem and the independence of the conditions of adjointness in the case of a curve for which the tangents at the multiple points are distinct from one another."

(11) Professor E. B. VAN VLECK: "A proof of the convergence of the Gaussian continued fraction for

$$\frac{F(a, \beta+1, \gamma+1, x)}{F(a, \beta, \gamma, x)}."$$

(12) Professor T. E. McKINNEY: "Some new kinds of continued fractions."

(13) Professor E. D. ROE: "Note on symmetric functions."

(14) Dr. G. A. MILLER: "Groups defined by the orders of two generators and the order of their product."

(15) Dr. H. F. STECKER: "On the determination of surfaces capable of conformal representation upon the plane so that the geodetic lines shall be represented by a prescribed system of plane curves."

(16) Dr. C. N. HASKINS: "On the invariants of quadratic differential forms."

(17) Dr. EDWARD KASNER: "The cogredient and digredient theories of double binary forms."

(18) Professor MAXIME BÔCHER: "On Wronskians of functions of a real variable."

(19) Mr. F. G. RADELFINGER: "The analytical representation of a multiform function in the domain of an isolated singular point."

(20) Dr. VIRGIL SNYDER: "On the forms of unicursal sextic scrolls with a multiple linear directrix and one double curve."

(21) Dr. H. F. STECKER: "Concerning the osculating plane of m-fold space filling curves of the Hilbert-Moore type."

(22) Dr. H. F. STECKER: "On non-euclidean properties of plane cubics and of their first and second polars."

(23) Professor L. E. DICKSON: "Theory of linear groups in an arbitrary field."

(24) Mr. H. L. RIETZ: "On primitive groups of odd composite order."

(25) Miss I. M. SCHOTTENFELS: "On the non-isomorphism of two simple groups of order $8!/2$."

(26) Professor L. W. DOWLING: "On the generation of plane curves, of any order higher than four, with four double points."

(27) Professor L. E. DICKSON: "The configuration of the 27 lines on a cubic surface and the 28 bitangents to a quartic curve."

(28) Professor E. H. MOORE: "Concerning the second mean value theorem of the integral calculus."

(29) Mr. I. E. RABINOVITCH: "The application of circulants to the solution of algebraic equations."

(30) M. EMILE LEMOINE: "Note sur la construction approchée de π de Mr. George Peirce."

(31) Dr. C. W. McG. BLACK: "The parametric repre-

sentation of the neighborhood of a singular point of an analytic surface."

(32) Professor ALEXANDER PELL: "Some remarks on surfaces whose first and second fundamental forms are the second and first respectively of another surface."

Professor Pringsheim's papers were presented to the Society through Professor Moore, Dr. Haskins's through Professor Osgood, Mr. Rietz's through Dr. Miller, Dr. Rabinovitch's through Professor Morley, M. Lemoine's through the Secretary. In the absence of the authors the first paper of Professor Bôcher and the papers of Dr. Haskins and Dr. Black were read by Professor Osgood, Professor Pringsheim's papers by Professor Moore, M. Lemoine's paper by Professor Brown. The second paper of Professor Bôcher, the papers of Mr. Rietz, Miss Schottenfels, and Professor Pell, and the second paper of Professor Moore were read by title.

In Professor Bôcher's first paper the following theorem is proved by making use of the Riccati's equations satisfied by the ratios of the functions $\varphi_2 y' - \varphi_1 y$ and $\psi_2 y' - \psi_1 y$: If, in a certain interval, p, q, φ_2, φ_1, ψ_2, ψ_1 are continuous real functions of the real variable x, and if the last four of these functions have continuous derivatives, then, y being a solution not identically zero of the differential equation

$$y'' + py' + qy = 0,$$

the roots of the functions

$$\varphi_2 y' - \varphi_1 y, \quad \psi_2 y' - \psi_1 y$$

will separate each other if no one of the three functions

$$\varphi_1\psi_2 - \varphi_2\psi_1, \quad \varphi_1'\varphi_2 - \varphi_1\varphi_2' + \varphi_1^2 + p\varphi_2\varphi_1 + q\varphi_2^2,$$
$$\psi_1'\psi_2 - \psi_1\psi_2' + \psi_1^2 + p\psi_2\psi_1 + q\psi_2^2$$

vanishes at any point of the interval in question.

Certain extensions of this theorem are established and some important special cases are considered. The relations of the subject to Sturm's paper in *Liouville's Journal*, volume 1 (particularly pp. 149–164), and to earlier papers of the author are discussed.

The paper by Dr. Hutchinson considers the functions which are automorphic for the monodromic group of the Riemann surface $w^2 = (z - x_1)(z - x_2)(z - \mu_1)^2(z - \mu_2)^2$. The moduli of periodicity of the integrals of the first kind

are expressible in terms of a single parameter ζ the group of transformations on which is generated by the two

$$S\colon\ \zeta' = \zeta + \sqrt{3}, \qquad T\colon\ \zeta' = -1/\zeta.$$

The automorphic functions of ζ have for natural boundary the axis of reals, and for a fundamental region one limited by the unit circle and two parallels to the imaginary axis at the distances $\pm \frac{1}{2}\sqrt{3}$. Every such function is rationally expressible in terms of any one of the Rosenhain moduli.

In his first paper Professor Pringsheim recurs to the important lemma of Goursat (*Transactions*, volume 1 (1900), No. 1) and gives in detail its application to the proof of Goursat's form of Cauchy's theorem. Of this application Goursat gave a brief indication, which was subject to misinterpretation. The present paper contains also critical remarks concerning the possible crinkliness of the curve of integration of the integral if the theorem holds by this proof, as well as a more general form of the lemma valid and available in some cases of greater crinkliness.

In his memoir "Über die Multiplication bedingt convergenter Reihen" (*Mathematische Annalen*, volume 21 (1883), p. 327) Professor Pringsheim pointed out that the product series resulting from the application of Cauchy's rule of multiplication to two conditionally convergent series or to a conditionally convergent and a divergent series may be be unconditionally convergent, and suggested the query whether the same conclusion might hold for the product series of two conditionally convergent series. Professor Cajori (*Transactions*, volume 2 (1901), No. 1) showed that this may happen for the product series of two conditionally convergent series and also for the product series of a conditionally convergent and a divergent series. In his second paper Professor Pringsheim proves that all these results and the similar result for the product series of two divergent series are precisely typical for large categories of cases which are obtained by extremely simple methods depending primarily upon the properties of a power series on its circle of convergence.

In studying the convergence of any one of the well-known developments for an arbitrary function $f(x)$ in terms of Bessel functions (developments such as occur for instance in the study of mathematical physics) it becomes desirable

to have some simple relation connecting $J_n(z)$ with the trigonometric functions, $J_n(z)$ being here used to represent Bessel's function of order n. Mr. Ford's paper in dealing with the relationship between the two classes of functions establishes two theorems which are of especial value in connection with the study of series developments such as have just been mentioned, though they may be used to advantage in any study relative to the behavior of $J_n(z)$ in the neighborhood of the point $z = \infty$.

In his first paper Professor Moore considers the various current types of improper simple definite integrals (connected with the names of (1) Cauchy, Riemann, du Bois-Reymond, Dini, Schoenflies; (2) Harnack, Jordan; (3) Hölder; (4) de la Vallée-Poussin) and, generalizing their diversities, defines a system of types, which differ according to the way in which the integral depends definitionally in its limiting process or processes upon the set of points of singularity of the integrand function with respect to definite integration. The properties of the general type of this system are investigated.

Professor Bolza's paper gives a simple proof of Professor Osgood's theorem on a characteristic property of a strong minimum in the calculus of variations (*Transactions*, July, 1901). The paper will be published in the *Transactions*.

The paper by Dr. Bliss concerns the problem of the calculus of variations in which one end point is restricted to lie upon a given curve while the other is fixed. Sufficient conditions for a maximum or minimum in this case are derived by Kneser in his Variationesrechnung and some of them are also proved to be necessary. The purpose of this paper was to find a complete set of necessary and sufficient conditions by the method which Weierstrass used for fixed end points. By this method the necessity of the conditions was proven for Kneser's exceptional case when the fixed end point lies beyond the "Brennpunkt," at which, besides, the envelope of the field has a multiple point of a particular kind. It cannot, however, be applied when "Brennpunkt" and end point are coincident. From the discussion it follows that the "Brennpunkt" always lies nearer to the given curve than the conjugate point.

Dr. Field's first paper is in abstract as follows: Let $F(z, u) = 0$ be the equation of an irreducible algebraic

curve of degree n. The multiple points of the curve we shall assume to be double points with distinct tangents. Further we shall assume that no tangent at a double point, no inflexional tangent, and no asymptote is parallel to either of the axes, and also that the asymptotes are all distinct from one another, and no two parallel to each other. The coefficient of u^n we shall take equal to 1. In § 2 of his paper "On the reduction of the general abelian integral,"* the author has given the formula

$$\frac{du}{dz} = \sum_{\lambda} \frac{\gamma_\lambda F(a_\lambda, u)}{(z - a_\lambda)(u - b_\lambda)},$$

where the summation extends to all branch and double points (a_λ, b_λ), u being regarded as dependent variable, and where

$$\gamma_\lambda = \frac{1}{[F''_{u}{}_{a_\lambda, b_\lambda}]} \quad \text{or} \quad \frac{2}{[F''_{u}{}_{a_\lambda, b_\lambda}]}$$

according as (a_λ, b_λ) is a branch point or a double point.

On reducing the above formula to the form

$$\begin{aligned}\frac{du}{dz} &= \sum_{\lambda} \frac{\gamma_\lambda F(a_\lambda, u)}{(z - a_\lambda)(u - b_\lambda)} \\ &= z^{-1} p_1(z, u) + z^{-2} p_2(z, u) + \cdots + z^{-k} p_k(z, u) + \cdots,\end{aligned}$$

where $p_1(z, u), p_2(z, u), \cdots$ are homogeneous polynomials in z, u of degree $n - 1$, we readily see that the coefficients of all the terms in the polynomials $p_1(z, u), \cdots, p_{n-2}(z, u)$ must vanish and that $z^{-(n-1)} p_{n-1}(z, u)$ must reduce to the single term $z^{-1}u$, so that we have

$$\frac{du}{dz} = z^{-1}u + z^{-n} p_n(z, u) + \cdots$$

The conditions implied upon the constants γ_λ by the vanishing of the coefficients here in question may be shown to be

$$\sum_{\lambda} \gamma_\lambda a_\lambda{}^i b_\lambda{}^k = 0 \qquad (i + k = 0, 1, 2, \cdots, n - 2),$$

with the exception of the one condition corresponding to the pair of values $i = 0$, $k = n - 2$, which takes the form

$$\sum_{\lambda} \gamma_\lambda b_\lambda{}^{n-2} = 1.$$

* *Trans. Amer. Math. Soc.*, vol. 2 (1901), pp. 49–86.

These conditions may be included in the single formula

$$\sum_{\lambda} \gamma_\lambda G(a_\lambda, b_\lambda) = d_{0, n-2}$$

where

$$G(z, u) = \sum d_{i,k} z^i u^k \qquad (i + k \leqq n - 2)$$

is an arbitrary polynomial in z, u of degree $n - 2$.

On substituting their values for the quantities γ_λ we obtain a formula which may conveniently be written in the form

$$\sum_{\lambda} \frac{G(a_\lambda, b_\lambda)}{[F''_{u}{}_{a_\lambda, b_\lambda}]} = d_{0, n-2}$$

on supposing an element of the summation corresponding to a double point to be repeated.

A similar formula is obtained on regarding z as the dependent variable.

Dr. Field's second paper is summarized as follows: Let $F(z, u) = 0$ be the equation to an irreducible algebraic curve of degree n. It is supposed that the tangents at the multiple points are distinct from one another and that these points are not at the same time branch points, and further it is assumed that the asymptotes are all distinct from one another, none of them parallel to the axis of u, and no two parallel to each other.

In a paper presented at the meeting of the Chicago Section in December last, the author proved the Riemann-Roch theorem and the independence of the conditions of adjointness in the case where the multiple points are all double points. The present paper extends the methods there employed to multiple points of higher order.

It is shown that the most general rational function which becomes infinite only at ∞ is given by the expression

$$\sum_{\lambda} \sum_{s=0}^{\sigma_\lambda - 2} \sum_{r=0}^{\sigma_\lambda - s - 2} c_{r,s,\lambda} \left(\frac{\partial}{\partial a_\lambda}\right)^r \left(\frac{\partial}{\partial b_\lambda}\right)^s \frac{F(a_\lambda, u)}{(z - a_\lambda)(u - b_\lambda)} + T(z, u),$$

where $T(z, u)$ is an arbitrary polynomial and the coefficients $c_{r,s,\lambda}$ arbitrary constants, where σ_λ is the order of the multiple point (a_λ, b_λ), and where the summation extends to all the multiple points (a_λ, b_λ). This may be written in the form

$$\sum_{\lambda} \gamma_\lambda \frac{F(a_\lambda, u)}{(z - a_\lambda)(u - b_\lambda)} + T(z, u)$$

on putting

$$\gamma_\lambda = \sum_{s=0}^{\sigma_\lambda - 2} \sum_{r=0}^{\sigma_\lambda - s - 2} c_{r,s,\lambda} \left(\frac{\partial}{\partial a_\lambda}\right)^r \left(\frac{\partial}{\partial b_\lambda}\right)^s.$$

On indicating by $(a_1, b_1), \cdots, (a_\delta, b_\delta)$ the multiple points, supposed to be δ in number, and by $(a_{\delta+1}, b_{\delta+1}), \cdots, (a_{\delta+Q}, b_{\delta+Q})$ certain Q other points, we consider the conditions under which the function

$$\sum_{\lambda=1}^{\delta+Q} \gamma_\lambda \cdot \frac{F(a_\lambda, u)}{(z - a_\lambda)(u - b_\lambda)} + T(z, u)$$

remains finite at ∞, an operator γ_λ here reducing to an arbitrary constant in case (a_λ, b_λ) is an ordinary point or a double point. We find that the polynomial $T(z, u)$ must reduce to an arbitrary constant c, and on expressing our function in the form

$$\sum_{\lambda=1}^{\delta+Q} \gamma_\lambda \cdot \frac{F(a_\lambda, u) - F(a_\lambda, b_\lambda)}{(z - a_\lambda)(u - b_\lambda)} + C$$

$$= z^{-1} p_1(z, u) + z^{-2} p_2(z, u) + \cdots,$$

where the functions $p_1(z, u), p_2(z, u), \cdots$ are homogenous polynomials in (z, u) of degree $n - 1$, we see that the coefficients must all vanish in the $p - 2$ polynomials $p_1(z, u), \cdots, p_{n-2}(z, u)$. This imposes certain conditions on the constants $C_{r,s,\lambda}$ involved in the operators γ_λ, and these conditions we find to be given by the equations

$$\sum_{\lambda=1}^{\delta+Q} \gamma_\lambda \cdot a_\lambda^i b_\lambda^k = 0 \qquad (i + k = 0, 1, 2, \cdots, n - 3).$$

An operator γ_λ operates throughout on any function (a_λ, b_λ) which it may happen to precede.

From the fact that a rational function which nowhere becomes infinite must be a constant, follows in the case where $Q = 0$ that the constants involved in the operators γ_λ must all have the value 0, and the interpretation of this necessity proves the independence of the conditions of adjointness. In the case $Q > 0$ our equations show that the number of arbitrary constants $C_{r,s,\lambda}$ involved in their solutions is $Q - q$, where q is the strength of the set of Q points $(a_{\delta+1}, b_{\delta+1}), \cdots, (a_{\delta+Q}, b_{\delta+Q})$ in determining an adjoint polynomial of degree $n - 3$, and from this the Riemann-Roch theorem immediately follows.

Professor Van Vleck considered the convergence of the well-known continued fraction of Gauss for

$$G(a, \beta, \gamma. x) = \frac{F(a, \beta + 1, \gamma + 1, x)}{F(a, \beta, \gamma, x)}$$

This has been previously shown by Thomè and Riemann to converge to $G(a, \beta, \gamma, x)$ as its limit over the entire complex plane with the exception of the portion of the real axis included between $x = 1$ and $x = +\infty$. The proofs of both of these writers are, however, of a special and somewhat involved character, not based upon considerations which can be applied to prove the convergence of many other continued fractions. The object of the present paper was to demonstrate the convergence by principles of a fundamental character for the theory of algebraic continued fractions. For this purpose two new theorems are established, the first of which is as follows:

I. If in a continued fraction

$$\frac{a_1 x}{1} + \frac{a_2 x}{1} + \frac{a_3 x}{1} + \cdots$$

the greatest modulus of any point of condensation of the sequence $a_1, a_2, a_3, \ldots$ is k, then within a circle of radius $\frac{1}{4k}$, described about the origin as center, the continued fraction will represent an analytic function, and the only singularities of the function contained within the circle will be poles. In any region excluding these poles and lying in the interior of the circle the convergence will be uniform.

This theorem will be further developed and published in a separate paper. The other theorem is as follows:

II. If from and after some fixed point in the continued fraction

$$\frac{1}{b_1 z} + \frac{1}{b_2} + \frac{1}{b_3 z} + \frac{1}{b_4} + \cdots$$

b_n is real and positive, and if $\sum b_n$ is divergent, the continued fraction will converge over the entire plane of z with the exception of 1° the whole or part of the negative half of the real axis, and 2° a series of isolated points $p_1, p_2, \cdots$ which lie without this half of the axis. The limit of the continued fraction is an analytic function whose only singularities exterior to the negative half axis are the points $p_1, p_2, \ldots$ which are poles.

By applying these theorems to the continued fraction of

Gauss, its convergence can be immediately demonstrated when a, β, γ are real. In case one or more of these three elements are complex, the second theorem must first be suitably modified and may then be employed as before. The two theorems, either separately or together, can be applied to all, or nearly all, the common algebraic continued fractions.

Professor McKinney discussed certain new kinds of continued fractions. These are characterized by 1° The system of intervals into which the series of real numbers is divided; and 2° the equation connecting two consecutive complete quotients. The paper defines these continued fractions and states some laws obeyed by developments representing the square root of integers, comparing these laws with the like cases of ordinary continued fractions.

Professor Roe's paper first calls attention to other proofs, and then gives a proof of Professor W. H. Metzler, that in the product of a symmetric function

$$\sum a_1^{p_1} a_2^{p_2} \cdots a_m^{p_m}$$

by the alternant $(0, 1, 2, \cdots, m-1)$ the result is obtained by adding the p's in all possible permutations to the exponents of the columns of the alternant written as a determinant, in which each line contains the powers of a single letter, thus giving the product in the general case as the sum of $m!$ alternants. This proof consists in combining the theorem for the multiplication of a determinant of the nth order by a polynomial of n terms, with the fact that every symmetric function is a function of the s's, each of which is a polynomial of n terms.

Attention is next called to the fact that in the preceding product the coefficient of the alternant $(i_1, i_2, \cdots, i_m)$ is equal to the coefficient of $a_{i_1} a_{i_2} \cdots a_{i_m}$ in the determinant

$$\begin{vmatrix} a_{i_1} & a_{i_2} \cdots & a_{i_m} \\ a_{i_1-1} a_{i_2-1} & \cdots & a_{i_m-1} \\ \cdots\cdots & \cdots\cdots & \cdots\cdots \\ a_{i_1-m+1} & \cdots & a_{i_m-m+1} \end{vmatrix}$$

which determinant is represented by $\{i_1 i_2 \cdots i_m\}$, and that a symmetric function $a_0^n \sum a_1^{p_1} a_2^{p_2} \cdots a_m^{p_m}$ is equal to a sum of determinants of the preceding form of the nth order in the

a's, with a similar coefficient excepting a sign factor, so that we obtain, with certain obvious details which need not be given here, the two formulas

$$(1)\quad (0, 1, 2, \cdots, m-1)\Sigma a_1^{p_1} a_2^{p_2} \cdots a_m^{p_m} = \Sigma \begin{Bmatrix} p_1 p_2 \cdots p_m \\ i_1 i_2 \cdots i_m \end{Bmatrix} (i_1 i_2 \cdots i_m),$$

$$(2)\quad a_0^n \Sigma a_1^{p_1} \cdots a_m^{p_m} = \Sigma (-1)^\mu \begin{Bmatrix} p_1 p_2 \cdots p_m \\ i_1 i_2 \cdots i_m \end{Bmatrix} \{ i_{m+1} i_{m+2} \cdots i_{m+n} \}.$$

Let l, m, n represent the orders of three operators L, M, N respectively and let $LM = N$. It is known that the group generated by any two of these operators is completely defined by the values of l, m, n whenever one of the following conditions is satisfied: 1° Two of these numbers are equal to 2; 2° one of them equals 2 the other 3, while the third has one of the three values, 3, 4, 5. The object of Dr. Miller's paper is to prove that two of the operators mentioned above can be so selected as to generate any one of an infinite system of groups of finite order whenever the values of l, m, n do not satisfy one of the two given conditions. This theorem is proved by observing that substitutions corresponding to L and M can be so selected as to generate a transitive group whose degree is an arbitrary multiple of some finite number.

Dr. Stecker's first paper is in continuation of a paper by the same author under nearly the same title in the *Transactions*, volume 2, p. 152. It is shown that the problem depends upon the integration of the system of partial differential equations

$$(1)\quad \begin{aligned} EF_u - \tfrac{1}{2}EE_v - \tfrac{1}{2}FE_u &= \frac{a_1}{a_5}\Delta, \\ -GF_v + \tfrac{1}{2}GG_u + \tfrac{1}{2}FG_v &= \frac{a_4}{a_5}\Delta, \\ EG_u - \tfrac{3}{2}FE_v - \tfrac{1}{2}GE_u + FF_u &= \frac{a_2}{a_5}\Delta, \\ -GE_v + \tfrac{3}{2}FG_u + \tfrac{1}{2}EG_v - FF_v &= \frac{a_3}{a_5}\Delta, \end{aligned} \qquad \left(E_u = \frac{\partial E}{\partial u}, \text{ etc.}\right)$$

where a_i has the same meaning as in the former paper and $\Delta = EG - F^2$. These lead to two relations

$$\frac{\Delta}{\varphi(v)E^{\frac{1}{2}}} = e^{\int \frac{a_2}{a_5} du} \quad e^{-3\int \frac{a_1}{a_5}\frac{F}{E} du},$$

$$\frac{G^{\frac{1}{2}}}{\varphi(u)\Delta} = e^{\int \frac{a_3}{a_5} dv} \quad e^{-3\int \frac{a_4}{a_5}\frac{F}{G} dv}$$

The paper considers the integration of system (1) for $a_1 = a_4 = 0$ and the conclusion is that there are two solutions corresponding to one real and one imaginary surface respectively; that a real solution requires that a_2 and a_4 also vanish.

The problem of determining the number of the differential invariants of quadratic differential forms has been attacked by Zorawski and Levi-Civita. Both of these investigators used the methods of Lie's theory of continuous groups. Zorawski, however, restricted his investigations to the binary forms, and Levi-Civita determined, not the precise number of the invariants, but a lower limit for that number. In Dr. Haskins's paper the determination of the exact number of invariants of the general form in n variables is carried out. The methods used are those of Lie's theory. The work proceeds mainly by mathematical induction, and in it much use is made of certain special forms which at some stages of the process can be used instead of the general form. The four-index symbols of Christoffel also play an important part in the demonstrations. The results are that for

$$\left.\begin{matrix} n \gtreqless 3, \\ \mu > 2, \end{matrix}\right\} \quad \text{or} \quad \left\{\begin{matrix} n = 2, \\ \mu \gtreqless 4, \end{matrix}\right.$$

there are

$$I_{n\mu} \equiv \frac{n(\mu-1)}{2} \frac{(n+\mu-1)!}{(n-2)!\,(\mu+1)!}$$

invariants of order μ for the form in n variables.

For $n \gtreqless 3$, $\mu = 2$ there are

$$J_{n2} \equiv \frac{(n-2)(n-1)n(n+3}{12}$$

invariants, and in the cases of the invariants of lowest orders for the binary forms, viz., $n = 2$, $\mu = 2$ and $\mu = 3$, there is a single invariant. The results for $n = 2$ agree with those of Zorawski. The methods used are, however, less complicated than his.

The methods employed in this paper can be applied to the determination of the number of simultaneous invariants of several forms. The results are that for m forms the number of simultaneous invariants of order μ, exclusive of the invariants of the separate forms, is, if $\mu = 0$,

$$\frac{mn(n+1)}{2} - n^2;$$

if $n = 2$, $\mu = 3$, or if $\mu = 2$, $n > 2$, it is

$$\frac{(m-1)n(n+\mu)!}{(\mu+1)!(n-1)!} + \frac{mn(n-1)}{2};$$

and in all other cases it is

$$I_{mn\mu} \equiv \frac{(m-1)n(n+\mu)!}{(\mu+1)!(n-1)!}.$$

In these last cases the invariants may be taken as simultaneous invariants of $m-1$ pairs of forms.

Forms containing two sets of binary variables $x_1 : x_2$ and $y_1 : y_2$ have been studied from two points of view. In what may be termed the cogredient theory, the variables are supposed to undergo the same linear transformation, so that the group defining concomitants is three-parametric

$$G_3 : \quad \begin{aligned} x_1 : x_2 &= r_1x_1' + r_2x_2' : r_3x_1' + r_4x_2', \\ y_1 : y_2 &= r_1y_1' + r_2y_2' : r_3y_1' + r_4y_2'. \end{aligned}$$

In the digredient theory, on the other hand, the variables are supposed to undergo independent linear transformations, so that the fundamental group is six-parametric

$$G_6 : \quad \begin{aligned} x_1 : x_2 &= r_1x_1' + r_2x_2' : r_3x_1' + r_4x_2', \\ y_1 : y_2 &= s_1y_1' + s_2y_2' : s_3y_1' + s_4y_2'. \end{aligned}$$

While the digredient concomitants of any forms are necessarily also cogredient concomitants of the same forms, the converse does not hold. After remarking that G_3 is a subgroup of G_6, consisting of those transformations of the latter which transform the bilinear form

$$\sigma \equiv x_1y_2 - x_2y_1$$

into itself, Dr. Kasner establishes the following relation between the two theories: The cogredient concomitants of any system S of double binary forms coincide with the di-

gredient concomitants of the enlarged system S, σ obtained by adjoining σ to the original system. This is then generalized so as to apply to the group transforming an arbitrary bilinear form into itself. The results are interpreted in the theory of correspondences, in the geometry of reciprocal radius vectors, and in the geometry on the quadric surface. Special applications are made to systems of bilinear forms and to the quadri-quadric form. In the latter case use is made of the connection between double binary and quaternary forms.

In his note in the 23d volume of the *Acta Mathematica*, Mittag-Leffler has greatly extended the domain in which a function can be represented by a single analytic expression. But his results are restricted to uniform functions or to one branch of a multiform function, since the independent variable is restrained from turning about singular points. Mr. Radelfinger's paper concerns itself with removing this latter restriction and constructing an expression uniformly convergent within a multiform domain surrounding an isolated singular point, a branch point, and containing no other singular point. The method employed is founded on the notion of analytic prolongation and is briefly as follows:

A branch point of a function $F^{(z)}$ is chosen as origin 0 and a small circumference r_0 drawn around it passing through all the leaves connected at 0. A non-crossing curve C passing between 0 and all other singular points is next drawn. The curve is constructed to enclose an area of maximum extent and traverses all the leaves. It may be closed or not. The region included between r_0 and C is denoted by S. A fixed point a is then located in the region S and any other point Z taken. A curve C is drawn connecting a and Z and lying wholly within S. It is then shown that for a chosen unique initial value $F_k(a)$ the function $F_k(z)$ remains uniform within S. A length ρ is then chosen equal to the minimum distance of any point C from the boundary of S and the arc az divided into n parts locating $n-1$ intermediate points ξ such that

$$|z-\xi_{n-1}| = |\xi_{n-1}-\xi_{n-2}| = \cdots |\xi_1 - a| < a^n\rho \quad (0<a<1).$$

Beginning with ξ_{n-1}, circles are described about the points ξ and the point a, the radii of the circles decreasing uniformly from ρ which surrounds ξ_{n-1} to $a^n\rho$ which surrounds a. Starting with the Taylor's expansion for $F_k(z)$ in the domain ξ_{n-1}, the function is prolonged backward along c,

and the successive derivatives $F^{\lambda}(\xi_m)$ are expressed in terms of the derivatives $F(\xi_{m-1}^{\lambda_1 + \lambda_2})$, until a is reached. A series n times infinite is thereby obtained which is proved to converge uniformly for all points within S for a less than one and n sufficiently large. The complete demonstration is not given but it is carried up to a point beyond which Mittag-Leffler's analysis applies without change. An interesting simplified form of the expression is obtained for the case where a circle R can be substituted for the curve C without excluding Z from the region S. To deduce this, a circle of radius r is substituted for C, and a taken as (r, a). By choosing Z sufficiently near the origin, so that ρ can be taken equal to ar, it is shown that in order to make a circuit of the origin n must be taken greater than 6.

In Dr. Snyder's paper those ruled surfaces are studied which can be generated by the line of intersection of two developables, the sum of whose classes is six. The double curve is in general of order 10, but this may be composite: only those forms are considered in which the components are straight lines and a non-reducible curve. Twenty-eight types are found.

In Dr. Stecker's second paper the broken curves C_n, of which the curve is the limit, are given for the three dimensional Hilbert curve and it is shown by considering secant planes that the osculating plane to curves of this general type are everywhere indeterminate.

Dr. Stecker's third paper is in continuation of a paper by the author in the *American Journal of Mathematics*, volume 22, p. 31, and is concerned with the properties which plane cubics and their polars have in the hyperbolic and elliptic geometries. The paper will appear in the *American Journal*.

The object of several recent papers by Professor Dickson is to show that various branches of group theory may be correlated by a treatment of groups of transformations in a given domain of rationality. The first of these memoirs, to appear in the *Transactions* for October, 1901, presents the fundamental conceptions of the theory and sets up four infinite systems of groups of transformations which are simply groups in every domain of rationality. For the case of the field of all complex numbers, these groups are the simple continuous groups of Sophus Lie. Aside from the isolated

groups, the latter give the only systems of simple continuous groups of a finite number of parameters. Corresponding to the isolated group of 14 parameters, there exists in the Galois field of order p^n a (new) system of simple groups of order $p^{6m}(p^{6m}-1)(p^{2m}-1)$.

The second memoir, "Linear groups in an infinite field," communicated to the London Mathematical Society, June 13, 1901, investigates the linear group in an arbitrary field F which leaves invariant

$$\xi_0^2 - \nu\eta_0^2 + \xi_1\eta_1 + \xi_2\eta_2 + \cdots + \xi_m\eta_m,$$

where ν is not square in F, thus obtaining new simple groups. Let Q be the field obtained by the extension of F by a root of $x^2 = \nu$ and let $\bar{y}$ denote the conjugate of y with respect to F. The linear transformations on $\xi_i, \eta_i (i = 1, \cdots, m)$, with coefficients in Q of determinant unity, which leave invariant

$$\sum_{i=1}^{m} (\xi_i\bar{\eta}_i - \bar{\xi}_i\eta_i)$$

form the hyperabelian group. Its maximal invariant subgroup is formed of the transformations

$$\xi_i' = \mu\xi_i, \quad \eta_i' = \mu\eta_i \quad [\mu\bar{\mu} = 1, \mu^{2m} = 1] \quad (i = 1, \cdots, m).$$

Hence the group of linear fractional hyperabelian transformations of determinant unity is simple in every field.

The third memoir, offered July 18, 1901, to the *Quarterly Journal of Mathematics*, investigates the group in an arbitrary field which corresponds to the isolated simple continuous group of 78 parameters.

Mr. Rietz's paper is in abstract as follows: Let G be any non-regular transitive group of order g and degree n. Let G_1 be the subgroup of G containing all the substitutions leaving a given letter fixed. It is proved that G then contains $> \frac{g}{x+m}$ substitutions of degree $< n$, where x is the number of transitive constituents of G_1 and m the number of elements of G not occurring in G_1. When G is primitive, $m = 1$, so that a primitive group of degree n and of order g contains more than $\frac{g}{x+1}$ substitutions of degree $< n$. In particular, in a multiply transitive group of degree n more than one-half the substitutions are of degree $< n$.

From the above the following theorem is readily obtained: If G is a primitive group of degree kp (p a prime) and of order mp (m prime to p and $p-1$) the subgroup G_1 has at least $p+1$ systems of intransitivity.

In a primitive group G of odd order, G_1 is intransitive. Some theorems are given restricting the order of G_1 when it has a transitive constituent of given form. Thus, if G_1 has a transitive constituent of prime degree p, its order is not divisible by p^2.

Nearly all the theorems given have some bearing on the problem of determining all the primitive groups of odd order of a given degree. The results of a determination of all these groups of degree < 150 show that, aside from invariant subgroups of the metacyclic groups, there are only 8 such groups of degree < 150. Their orders are the following numbers: 25.3, 27.13, 27.39, 81.5, 121.3, 121.15, 125.31, 125.31.3. The first factor in these numbers as they are written gives the degree of the group of that order. Each of these groups is solvable.

In a paper by Timerding (*Crelle*, volume 123) it is shown that plane quintics with four double points may be generated by the intersection of corresponding elements of two projectively related forms, one being a linear pencil of conics and the other a quadratic pencil, viz., the system of tangents to a conic.

This theorem is easily proven in various ways and can be generalized to inelude the generation of any curve having four double points.

In a paper read before the Toronto Meeting of the Society, summer of 1897, Professor Dowling showed that the equation of any plane curve of order n may be put into the form

$$\alpha\beta U + \varphi^2 \Pi = 0$$

in a variety of ways, where α and β are linear expressions in the variables, U is an expression of order $n-2$, φ is a quadratic, and Π consists of the continued product of $n-4$ linear factors.

This curve will have four double points if U is made to pass through the four intersections of $\alpha\beta = 0$ and $\varphi = 0$. U will then be of the form $U \equiv \alpha\beta P + 2\varphi Q$, where P and Q are each expressions of order $n-4$. The curve will then have for its equation $\alpha^2\beta^2 + 2\varphi Q\alpha\beta + \varphi^2\Pi = 0$ and is thus evidently generated by the two projectively related forms

$$\lambda^2 P + 2\lambda Q + \Pi = 0$$

and

$$\alpha\beta - \lambda\varphi = 0.$$

The first of the forms is a quadratic pencil of curves of order $n-4$ characterized by being tangent to the curve

$$PH - Q^2 = 0.$$

This latter curve is of order $2n-8$ and possesses $n-4$ multiple tangents each touching the curve in $n-4$ points.

In Professor Dickson's second paper it is shown that the cubic form C in 27 variables with 45 terms, which was given by Cartan in his thesis on simple continuous groups, is suitable to define the configuration of the 45 triangles formed by the 27 straight lines on a general cubic surface.

In a memoir sent to the *Quarterly Journal of Mathematics*, July 18, 1901, the author studies the linear group in an arbitrary field which leaves invariant the cubic form C. For the case of a continuous field, the results agree with those of Cartan on the simple continuous group of 78 parameters. For the Galois field of order p^n, the corresponding group has the order

$$p^{36n}(p^{12n}-1)(p^{9n}-1)(p^{8n}-1)(p^{6n}-1)(p^{2n}-1)$$
$$(p^n-1)(p^n+2).$$

It may be represented as a transitive substitution group on $(p^{9n}-1)(p^{8n}+p^{4n}+1)$ letters.

The present paper also investigates the group of a quartic form Q in 56 variables with 630 terms. The partial derivative of Q with respect to any one of the variables is a cubic form differing from the above form C only in notation. The function Q defines the configuration of the 56 points of contact of the 28 bitangents to a quartic curve without double points.

In Professor Moore's second paper, from the second mean value theorem in the usual form

$$\int_a^b f(x)\varphi(x)dx = a\int_a^\xi \varphi(x)dx + \beta\int_\xi^b \varphi(x)dx,$$

$$(a = f(a) \leqq f(b) = \beta, \qquad a \leqq \xi = b),$$

where on the interval ab $f(x)$ is monotonic not decreasing and $\varphi(x)$ is integrable, is first deduced the theorem in the more general form

$$(a \leqq f(a+0) \leqq f(b-0) \leqq \beta, \qquad a \leqq \xi \leqq b)$$

stated by du Bois-Reymond (*Zeitschrift für Mathematik und Physik*, volume 20 (1875), p. 126) in his review of Thomae's book on definite integrals. Two simple deductions are given, the first being doubtless that which du Bois-Reymond had in mind. The general theorem is not well known, the only preceding proof being that of Pringsheim (*Münchener Berichte*, 1900) based on a suitable generalization of a proof of the simpler theorem.

The general theorem is then extended to the case of the general improper integrals defined in the author's first paper, by a generalization of Jordan's theorem (Cours d'analyse, 2d edition, volume 2 (1894), p. 223) of the existence of the integral on the left, and by a generalization of the method used by Harnack in proving the simpler theorem for the case of his improper integrals. A corollary of this theorem is the generalization of the modified first mean value theorem (du Bois-Reymond, *Mathematische Annalen*, volume 7 (1874), p. 605).

Since the times of Euler and Lagrange it is well known that the roots of an algebraic equation of the nth degree with one unknown, may be assumed to be linear functions of n variables (or $n-1$ variables, in case of an equation freed of the second term), in which the coefficients are different nth roots of unity. Dr. Rabinovitch in his paper remarks that these forms might yield an elegant solution in case of the quadratic, cubic, and quartic, on equating the elementary symmetric functions of these expressions of the roots to the corresponding coefficients. Thus in the case of the quartic, for instance, with second term eliminated, assuming the roots to be of the form

$$x_1 = y + z + t, \qquad x_2 = y - z - t,$$
$$x_3 = -y + z - t, \qquad x_4 = -y - z + t,$$

which is obtained from the form referred to above by putting

$$y + t = y' + t', \quad y - t = i(y' - t), \quad z = z',$$

we find their sum $= 0$, the sum of their products by twos $= -2(y^2 + z^2 + t^2)$, which is equated to the coefficient of x^2, and so on. Then he shows that the same result is obtained almost automatically, with a minimum of calculation, by equating the given equation to a circulant of corresponding order in which the leading or diagonal constituent x is the unknown of the given equation and the remaining constituents are the facients in the linear expressions of the roots.

Each of the given equations then becomes identical with a given circulant which represents the product of all the linear factors of the given equation. He establishes the generality of this procedure (with the exception, of course, of the possibility of determining the values of these facients by radicals from the obtained relations) by a theorem of Glaisher, according to which

$$C(a_0 a_1 a_2 \cdots a_{n-1}) = \prod_{r=0}^{r=n-1} (a_0 + a_1\omega_r + a_2\omega_r^2 + \cdots + a_{n-1}\omega_r^{n-1}).$$

(See Muir, Theory of Determinants § 149.)

In the case of the general quintic the same procedure yields in an easy manner a set of equations very similar to to the equations which McClintock, in his paper "Analysis of the quintic equation" (*American Journal of Mathematics*, volume 8), gives under the name of the Eulerian equations The first set may be obtained also, as indicated above, from the elementary symmetric functions of the linear expression for the roots. This last procedure, however, is very lengthy and laborious.

Dr. Black's paper, which will appear in the *Proceedings of the American Academy of Arts and Sciences*, volume 37, is in abstract as follows: Let $F(x, y, z) = 0$ be the equation of a surface, F being a function of the three independent variables x, y, z analytic in the point $x = a$, $y = b$, $z = c$; and suppose that the function and all its first partial derivatives vanish in this point. Then it is possible to represent the coördinates of all points of the surface $F = 0$ that lie in the neighborhood of the point (a, b, c) by means of a finite number of sets of parametric formulas, each of the type

$$x = f(u, v), \quad y = \varphi(u, v), \quad z = \psi(u, v),$$

where f, φ, ψ are functions of the parameters u, v, analytic each in the point $u = u_0$, $v = v_0$. A proof of this theorem was attempted by G. Kobb, *Journal de mathématiques*, series 4, volume 8 (1892), but his analysis is incomplete in essential points.

By means of a quadratic transformation of the type

$$x - a = x'(z - c), \quad y - b = y'(z - c) \qquad (1)$$

the neighborhood of the point (a, b, c), in general singular, is mapped upon the neighborhood of a curve upon a new surface. If this curve does not contain multiple factors, its neighborhood, except for a finite number of neighborhoods

of singular points first determined, is represented by a finite number of formulas, in each of which one coördinate is expressed as an analytic function of the other two. If the curve does, however, contain multiple factors the introduction of a quadratic transformation of the type

$$x = x'z \qquad (2)$$

(preceded by a certain other transformation of the form $x + py = x_1$) secures the same result as in the other case.

Each of the new singular points is then subjected to the same treatment as the original point, and the process is repeated until all the singular points disappear, the parametric representation then becoming at once possible. It is shown that this result will be reached by means of a finite number of transformations of types (1) and (2), together with certain other one to one transformations.

F. N. COLE.

COLUMBIA UNIVERSITY.

THE ITHACA COLLOQUIUM.

THE Third Colloquium of the AMERICAN MATHEMATICAL SOCIETY was held at Cornell University, Ithaca, N. Y., beginning on Wednesday, August 21, 1901, and extending over the following three days. Before describing the proceedings, it may be of interest to recall the work of the previous colloquia.

The first colloquium * organized by the Society was held in connection with its third summer meeting at Buffalo, N. Y., September 2–5, 1896. Two courses of six lectures each were delivered before an audience of thirteen members. Professor Maxime Bôcher discussed "Linear differential equations and their applications," and Professor James Pierpont, "The Galois theory of equations." The innovation proved so successful that the participants recommended to the Council that the same plan should be adopted the following summer; but the meeting at Toronto, with

* See the report, including abstracts of the courses of lectures, by Professor T. S. Fiske in BULLETIN, volume 3, pp. 49–59. Professor Bôcher's lectures were in part reproduced in the *Annals of Mathematics*, 1st series, vol. 12 (1898), pp. 45–53. Professor Pierpont's lectures were published in the same journal, 2d series, vol. 1 (1899), pp. 113–143, and vol. 2 (1900), pp. 22–56.

the British association for the advancement of science, rendered this impracticable. The second colloquium* was therefore held, after an interval of two years, in connection with the fifth summer meeting, at Cambridge, Mass., August 22–27, 1898. Professor W. F. Osgood delivered six lectures on "Selected topics in the general theory of functions," and Professor A. G. Webster six lectures' on "The partial differential equations connected with wave propagation." Twenty-six members attended the two courses.

On account of the intervening Paris exposition a period of three years elapsed between the second and third colloquia. At the December meeting, 1900, the Council appointed a committee, consisting of Professors F. N. Cole, W. F. Osgood, James Pierpont, J. H. Tanner, and H. S. White, to arrange for a third colloquium in connection with the eighth summer meeting. A preliminary notice containing the plans agreed upon was issued in May, 1901. In accordance with these plans, the following twenty-five members of the Society assembled Wednesday morning, August 21, 1901, in one of the lecture rooms of White Hall, Cornell University, and registered for the colloquium:

Dr. G. E. Bliss, Professor Oskar Bolza, Professor E. W. Brown. Professor F. N. Cole, Professor L. L. Conant, Professor T. S. Fiske, Mr. W. B. Ford, Mr. B. F. Groat, Dr. E. R. Hedrick, Professor T. F. Holgate, Dr. J. I. Hutchinson, Dr. Edward Kasner, Dr. G. H. Ling, Professor J. L. Markley, Professor E. H. Moore, Professor G. D. Olds, Professor W. F. Osgood, Dr. I. E. Rabinowitch, Mr. F. G. Radelfinger, Professor E. R. Skinner, Dr. Virgil Snyder, Professor J. H. Tanner, Professor E. B. Van Vleck, Professor L. A. Wait, Professor Alexander Ziwet.

The Society has always been fortunate in securing for the colloquia the services of lecturers of high standing. These serve purely as a matter of honor, receiving no compensation, except a very moderate allowance for their travelling and other expenses provided for by the collection of a nominal fee from each auditor.

During the four days of the colloquium, two courses of four lectures each were delivered by Professor Oskar Bolza, of the University of Chicago, and Professor E. W. Brown, of Haverford College. The titles and outlines of the courses are as follows:

*See the report by Professor H. S. White, BULLETIN, volume 5, pp. 57–58. Professor Osgood's lectures were published in the same volume, pp. 59–87.

PROFESSOR BOLZA :—THE SIMPLEST TYPE OF PROBLEMS IN THE CALCULUS OF VARIATIONS.

The principal object of the course was to give a detailed account of the simplest type of problems in the calculus of variations, in its historical development, with special emphasis upon the contributions of Weierstrass and his followers. After a critical account of the older theory of Euler, Legendre, and Jacobi, the lecturer passed to Weierstrass's theory and to the contributions made since Weierstrass by Schwarz, Kneser, Osgood, and others. The course concluded with an exposition of Hilbert's existence theorem, and with an indication of the lacunæ which must be filled before the theory of even the simplest type of problems can be declared complete.

PROFESSOR BROWN : — MODERN METHODS OF TREATING DYNAMICAL PROBLEMS AND IN PARTICULAR THE PROBLEM OF THREE BODIES.

The object of this course was to set forth some of the later attempts to introduce more rigor into the methods of solving dynamical problems, mainly the researches of Poincaré. The course was chiefly descriptive, in showing the principles of the methods, the mathematical difficulties which arise, and the results which have been obtained. Amongst the subjects treated were the following : The various forms of the differential equations of dynamics ; the theorems of Bruns and Poincaré on the non-existence of integrals ; absolute and relative invariant integrals ; solution by infinite series, including divergent series ; periodic and asymptotic solutions ; stability and instability. No special knowledge of any one subject was assumed.

Two lectures were delivered daily, one in the morning and one in the afternoon, the hour varying to suit the convenience of the lecturers and auditors. Each lecture extended through two hours, the period being divided by a recess of ten minutes. Professor Bolza's course was given on the mornings of Wednesday and Saturday, and the afternoons of Thursday and Friday ; Professor Brown's at the other four sessions.

The librarian of the University kindly offered the facilities of both the reading room and the stack rooms to members of the Society. The literature bearing on the subjects discussed in the colloquium, including autograph notes of Weierstrass's lectures, and a bibliography of the calculus of variations prepared by Dr. J. I. Hutchinson, was collected

and arranged for the convenience of the auditors. Printed syllabi of both courses were distributed, so that every opportunity was offered for intelligent participation. At the close of the courses, the members present, by a rising vote, unanimously expressed their appreciation of the value of the lectures, and of the self-sacrificing labor of the lecturers.

Friday evening the participants of the colloquium dined together at the Glenwood Hotel, and afterwards enjoyed a cruise around Cayuga Lake. During the remaining evenings there were social gatherings at the Town and Gown Club, whose privileges were thrown open to members of the Society. The hospitality of Cornell University, and of the mathematical department in particular, deserves the most grateful acknowledgment.

Detailed reports of the courses, prepared by the lecturers themselves, will appear in later numbers of the BULLETIN.

EDWARD KASNER,
Assistant Secretary.

COLUMBIA UNIVERSITY.

UPON THE NON-ISOMORPHISM OF TWO SIMPLE GROUPS OF ORDER 8!/2.

BY MISS IDA M. SCHOTTENFELS.

(Read before the American Mathematical Society, August 20, 1901.)

1. *Introduction.*—The proof offered in this paper of the non-isomorphism of the ternary linear fractional group Galois field $[2^2]$ and the alternating group of degree eight is shorter, simpler and more direct than that presented by the author in the *Annals of Mathematics*, volume 1, No. 3, April, 1900.

In a paper read December 29, 1900, before the Chicago Section of the AMERICAN MATHEMATICAL SOCIETY, by making use of the present method involving the three conjugate sub-groups (1) G_{168}, (2) G_{168}', (3) G_{168}'', the identity was established between the ternary group $G.F.[2^2]$ and the literal substitution group of degree 21, $G_{8!/2}^{21}$.

2. *The Ternary Group.*—In the above mentioned paper the group was defined as a group of fractional matrices, access being had to the group by means of the following matrices:

* $$\begin{matrix} A \\ \left[\begin{matrix} 010 \\ 001 \\ 100 \end{matrix}\right] \end{matrix}, \quad \begin{matrix} B \\ \left[\begin{matrix} 010 \\ 001 \\ 110 \end{matrix}\right] \end{matrix}, \quad \begin{matrix} C \\ \left[\begin{matrix} 010 \\ 001 \\ 1\rho 0 \end{matrix}\right] \end{matrix}, \quad \begin{matrix} D \\ \left[\begin{matrix} 010 \\ 001 \\ 1\rho^2 0 \end{matrix}\right] \end{matrix}, \quad \begin{matrix} I \\ \left[\begin{matrix} 100 \\ 010 \\ 001 \end{matrix}\right] \end{matrix};$$

where

$$\{A^3 = B^7 = C^7 = D^7 = I\}.$$

In this group there exist at least three conjugate subgroups of order 168, say (1) G_{168}, (2) G_{168}', (3) G_{168}'', such that (1) is transformable into (2) and (3) by elements of period three which leave A invariant; these groups are holoedrically isomorphic to the literal substitution group upon seven letters G_{168}^{7} and are thus defined

(1) G_{168} : $\{A^3 = B^7 = (BA^2)^2 = (AB^2)^4 = I\}$,

(2) G_{168}' : $\{A^3 = C^7 = (CA^2)^2 = (AC^2)^4 = I\}$,

(3) G_{168}'' : $\{A^3 = D^7 = (DA^2)^2 = (AD^2)^4 = I\}$,

where the elements B, C possess the following properties

$$(B^2C)^5 = (BC)^4 = (CB^4)^2 = I.$$

3. *The Alternating Group.*—The alternating group of degree eight and order 8!/2 or 20160 has a simple subgroup of order 168 upon seven letters which may be defined thus

$$G_{168}^{7} : \quad \{a^3 = b^7 = (ba^2)^2 = (ab^2)^4 = i\},$$

where i is the identity element, and $a = (234)\ (567)$, $b = (1346752)$.

There exists a group of order 36 upon eight letters which leaves $a = (234)\ (567)$ invariant, which group G_{36}^{8}, *if the alternating group is identical with the ternary group*, must possess an element of period three which transforms $b = (1346752)$ into c $(c^7 = i)$ where b, c satisfy the relations

$$(b^3c)^5 = (bc)^4 = (cb^4)^2 = i\,;$$

but *no such element exists*, therefore the two above mentioned groups *are non-isomorphic.*

CHICAGO,
December, 1900.

---

* 0, 1, ρ and ρ^2 are the marks of the Galois field $[2^2]$.

CONCERNING SURFACES WHOSE FIRST AND SECOND FUNDAMENTAL FORMS ARE THE SECOND AND FIRST FUNDAMENTAL FORMS RESPECTIVELY OF ANOTHER SURFACE.

BY PROFESSOR ALEXANDER PELL.

(Read before the American Mathematical Society, August 20, 1901.)

In the July number of the BULLETIN Dr. Eisenhart treated these surfaces and reached the following conclusions: "The ruled surfaces defined by the equations

$$y + \mu x = \sqrt{1+\mu^2} + C_2\mu + C_1$$
$$z - ix\sqrt{1+\mu^2} = (\mu + C_1\sqrt{1+\mu^2} + C_2)\cdot\frac{1}{i}$$

are the only surfaces whose first and second fundamental forms can be taken for the second and first fundamental forms of a surface. Further the second surface is only the first to a translation près. And of these surfaces the only real one is the sphere of radius unity."

In fact the above equations represent the spheres of radius unity and nothing else. For we have

$$y + zi + (\mu + \sqrt{1+\mu^2})\,x = (c_1 + 1)(\mu + \sqrt{1+\mu^2}) + c_2 + c_3$$

$$y - zi + (\mu - \sqrt{1+\mu^2})\,x = (c_1 - 1)(\mu - \sqrt{1+\mu^2}) + c_2 - c_3$$

or

$$(y-c_2)+(zi-c_3) = -(x-c_1)(\mu+\sqrt{1+\mu^2})+(\mu+\sqrt{1+\mu^2})$$

$$(y-c_2)-(zi-c_3) = -(x-c_1)(\mu-\sqrt{1+\mu^2})-(\mu-\sqrt{1+\mu^2}),$$

i. e.,

$$(y-c_2)^2 + (z+c_3 i)^2 + (x-c_1)^2 = 1.$$

But there are two more surfaces of the kind, both imaginary, both spheres of the radii ω, ω^2, respectively, where $\omega = \sqrt[3]{1}$.

For let

$$x = R\sin u\cos v, \quad y = R\sin u\sin v, \quad z = R\cos u.$$

Then

$$E = R^2, \quad G = R^2 \sin^2 u, \quad F = 0,$$

$$D = R, \quad D' = 0, \quad D'' = R \sin^2 u.$$

If now (1) $R = \omega$,

$$E_\omega = \omega^2, \; G_\omega = \omega^2 \sin^2 u, \; D_\omega = \omega, \; D_\omega'' = \omega \sin^2 u;$$

and if (2) $R = \omega^2$,

$$E_{\omega^2} = \omega, \; G_{\omega^2} = \omega \sin^2 u, \; D_{\omega^2} = \omega^2, \; D_{\omega^2}'' = \omega^2 \sin^2 u.$$

No other surface of the kind exists. For, as is shown in the article referred to, for these surfaces (ρ_1 and ρ_2 being two principal radii of curvature)

$$\rho_1 = \rho_2 = \frac{E}{D} = \frac{G}{D''},$$

i. e.,

$$\frac{D}{E} = \frac{D''}{G} = \lambda.$$

Bianchi has shown (Differentialgeometrie, German translation, p. 93, footnote), that λ is constant, if D, D', D'', E, F, G are proportional.

Using the equations (1) and (9) of Dr. Eisenhart's article, we have

$$\lambda^2 \cdot \sqrt{EG} + \frac{\partial}{\partial u}\left[\frac{1}{\sqrt{E}} \cdot \frac{\frac{\partial G}{\partial u}}{2\sqrt{G}}\right] + \frac{\partial}{\partial v}\left[\frac{1}{\sqrt{G}} \cdot \frac{\frac{\partial E}{\partial v}}{2\sqrt{E}}\right] = 0,$$

$$\frac{\sqrt{EG}}{\lambda} + \frac{\partial}{\partial u}\left[\frac{1}{\sqrt{E}} \cdot \frac{\frac{\partial G}{\partial u}}{2\sqrt{G}}\right] + \frac{\partial}{\partial v}\left[\frac{1}{\sqrt{G}} \cdot \frac{\frac{\partial E}{\partial v}}{2\sqrt{E}}\right] = 0.$$

Hence, on subtraction,

$$\lambda^2 \sqrt{EG} = \frac{\sqrt{EG}}{\lambda}, \qquad \lambda^3 = 1,$$

i. e.,

$$\lambda = 1, \; \omega, \; \omega^2.$$

UNIVERSITY OF SOUTH DAKOTA.

NOTES.

THE tenth regular meeting of the Chicago Section of the AMERICAN MATHEMATICAL SOCIETY will be held at Northwestern University, Evanston, Ill., on Friday and Saturday, December 27–28 next. Titles and abstracts of papers to be read at this meeting should be in the hands of the Secretary of the Section not later than December 5.

AMONG the officers of the American association for the advancement of science elected at the Denver meeting are: president of the association, Professor ASAPH HALL, U. S. Navy, retired; president of Section A (mathematics and astronomy), Professor G. W. HOUGH, of Northwestern University; secretary of Section A, Professor E. S. CRAWLEY, of the University of Pennsylvania. The next summer meeting of the Association will be held at Pittsburg, Pa., June 28 to July 3, 1902.

AT a meeting of the Liverpool Mathematical Society held in University College on Wednesday, June 26, 1901, Professor F. S. CAREY in the chair, Professor SEBASTIAN SIRCOM was unanimously elected president for the ensuing year, Professor CAREY, the retiring president, becoming vice-president. A paper on "Möbius' barycentric calculus and Grassmann's calculus of extension" was read by Professor R. A. SAMPSON, who explained the two methods and showed the connection between them. Professor L. R. WILBERFORCE exhibited some apparatus, using his springs showing the transformation of longitudinal into rotational oscillation, and vice versa. He also exhibited and explained the principal oscillations of the apparatus.

THE annual meeting of the German mathematical society was held at Hamburg, September 22–28. The preliminary programme of papers to be presented is as follows: V. EBERHARD, Halle: "Contribution to the theory of equations"; W. EBERT, Kiel: "On a question in celestial mechanics"; F. ENGEL, Leipsic: "Higher differential quotients"; F. FOLIE, Grivegnée: "On Oppolzer's formulæ for nutation"; D. HILBERT, Göttingen: "On certain recent mathematical dissertations"; J. KOWALEWSKY, Leipsic: "Report on the theories of Sophus Lie"; R. v. LILIENTHAL, Münster: "The geometry of motion in its application to differential geometry"; A. MARCUSE, Berlin: "Recent developments in the determination of geographical positions"; F. KLEIN, Göttingen, F. MEYER, Koenigsberg, and E. WIECH-

ERT, Göttingen: "An account of the present condition of the Encyclopædia of the mathematical sciences"; F. SCHILLING, Göttingen: "New kinematical models in tooth gear and their relation to the theory of contact transformations; P. SCHOUTE, Groningen: "On the mobility of a nullsystem N_{n-1} in R_n"; H. SCHUBERT, Hamburg: "On the number of constants of the n-dimensional generalization of polyedra;" P. STÄCKEL, Kiel: "Report on the development of instruction in applied mathematics in the German universities"; E. STUDY, Greifswald: "A new branch of geometry"; ADLER, Prague: "Spherical representation of surfaces and its application in descriptive geometry"; C. CHARLIER, Lund: "The astronomical explanation of an ice age"; HALM, Edinburgh: "The relations of terrestrial magnetism to seismological events and their significance for practical and theoretical astronomy"; F. MEYER, Koenigsberg: "The extension of the theorems of Henrici and Ivory to surfaces of any order"; TORKA, Berlin: "On a new geometry of motion"; E. ZERMELO, Göttingen: "On a maximum problem in geometry," "On the theory of the shortest lines."

THE several universities below offer during the winter semester 1901–1902 courses in mathematics as follows:

UNIVERSITY OF BERLIN.—By Professor L. FUCHS: Hyperelliptic functions, three hours; Theory of functions, three hours; Seminar, three hours.—By Professor H. A. SCHWARZ: Analytical geometry, four hours; Conic sections, two hours; Theory of analytical functions, four hours; Colloquium, two hours; Seminar, two hours.—By Professor J. KNOBLAUCH: Differential calculus, four hours; Applications of elliptic functions, four hours; Exercises in differential calculus, one hour.—By Professor R. LEHMANN-FILHÉS: Integral calculus, four hours, with exercises, one hour.—By Professor G. FROBENIUS: Theory of determinants, four hours; Theory of numbers, four hours; seminar, three hours.—By Professor K. HENSEL: Applications of elliptic functions to algebra and the theory of numbers, three hours; Mechanics, four hours; Colloquium, one hour.—By Dr. E. LANDAU: Algebra, four hours; Theory of transcendental functions, one hour.

UNIVERSITY OF BONN.—By Professor L. HEFFTER: Infinitesimal calculus, II., four hours, with exercises, one hour; Theory of curves and surfaces, four hours, with exercises, one hour.—By Professor H. KORTUM: Analytical geometry, five hours; Determinants, two hours; Seminar, two hours.

—By Professor R. LIPSCHITZ: Partial differential equations of physics, four hours; seminar, two hours.

UNIVERSITY OF BRESLAU.—By Professor J. ROSANES: Analytical geometry of space, three hours; Elements of the theory of functions, three hours; Seminar, one hour.—By Professor R. STURM: Theory of geometrical relations, II., two hours; Differential geometry, three hours; Seminar, two hours.—By Dr. F. LONDON: Infinitesimal calculus, four hours, with exercises, one hour.

UNIVERSITY OF ERLANGEN.—By Professor P. GORDAN: Infinitesimal calculus, four hours; Theory of numbers, four hours; Seminar, three hours.—By Professor M. NÖTHER: Analytical geometry, four hours; Analytical mechanics, four hours; Seminar.

UNIVERSITY OF FREIBURG.—By Professor J. LÜROTH: Analytical geometry and differential calculus, five hours; Seminar.—By Professor L. STICKELBERGER: Algebraic analysis, three hours; Elliptic functions, four hours.—By Dr. A. LOEWY: Differential equations, three hours; Political arithmetic, three hours.—By Dr. E. REBMANN: Elements of descriptive geometry with exercises, three hours.

UNIVERSITY OF GIESSEN.—By Professor M. PASCH: Plane analytical geometry, II., three hours and a half; Calculus of variations, two hours; Seminar, one hour.—By Professor E. NETTO: Infinitesimal calculus, three hours; Analytical mechanics, three hours; Theory of functions, continuation, two hours; Seminar, one hour.—By Professor R. HAUSSNER: Theory of numbers, four hours; Descriptive geometry, with exercises, four hours and a half; Exercises in higher mathematics, one hour and a half.

UNIVERSITY OF GÖTTINGEN.—By Professor F. KLEIN: Mechanics of particles and rigid systems, four hours; Seminar, two hours.—By Professor D. HILBERT: Potential, four hours; Number concept and quadrature of circle, two hours; Exercises in the theory of functions in seminar, two hours.—By Professor M. BRENDEL: Geodesy, two hours; Introduction to computation of perturbations, two hours; Elements of infinitesimal calculus, two hours.—By Professor F. SCHILLING: Descriptive geometry, four hours, with exercises, four hours; Kinematics, one hour.—By Professor K. BOHLMANN: Mathematics of insurance, three hours, with exercises in seminar, two hours.—By Dr. E. ZERMELO: Theory of numbers, four hours.—By Dr. J. SOMMER: Integral calculus, four hours.

UNIVERSITY OF GREIFSWALD.—By Professor W. THOMÉ: Theory of analytical functions, in particular elliptic functions, I., four hours; Infinitesimal calculus, I., four hours; Seminar, two hours.—By Professor E. STUDY: Analytical geometry, II., four hours; Differential equations, three hours; Seminar, one hour.

UNIVERSITY OF HALLE-WITTENBERG.—By Professor G. CANTOR: Theory of elliptic functions, two hours; Seminar, one hour.—By Professor A. WANGERIN: Analytical mechanics, four hours; Synthetic geometry, four hours, with exercises, one hour; Seminar, one hour.—By Professor V. EBERHARD: Analytical geometry of space, three hours, with exercises, one hour.—By Dr. H. GRASSMANN: Selected chapters of technical mechanics, two hours; Conformal representation, two hours, with exercises, one hour fortnightly.—By Dr. E. NEUMANN: Infinitesimal calculus, II., with exercises, four hours.

UNIVERSITY OF HEIDELBERG.—By Professor L. KOENIGSBERGER: Analytical mechanics, four hours; Selected chapters of integral calculus (calculus of variations, differential equations), two hours; Elliptic functions, two hours; Seminar, two hours.—By Professor H. VALENTINER: Theory of orbits, three hours.—By Professor M. CANTOR: Infinitesimal calculus, four hours, with exercises, one hour; Political arithmetic, two hours.—By Professor F. EISENLOHR: Optics, four hours; Infinitesimal calculus, five hours; Potential, two hours.—By Professor K. KOEHLER: Analytical geometry of space, three hours.—By Professor G. LANDSBERG: Descriptive geometry, with exercises, four hours.—By Dr. K. BOEHM: Partial differential equations, two hours; Abel's memoir on binomial series, one hour.

UNIVERSITY OF INNSBRUCK.—By Professor O. STOLZ: Analysis of real numbers, five hours; Seminar, two hours.—By Professor W. WIRTINGER: Algebraic functions and their integrals, five hours; Seminar, two hours.—By Dr. K. ZINDLER: Differential equations, three hours; Introduction to continuous groups, two hours.

UNIVERSITY OF JENA.—By Professor J. THOMAE: Analytical geometry of space, four hours; Elementary theory of functions, four hours; Seminar, two hours.—By Professor A. GUTZMER: Theory and applications of determinants, three hours; Integral calculus, four hours, with exercises, one hour.—By Professor J. FREGE: Theory of numbers, four hours.

UNIVERSITY OF KIEL.—By Professor L. POCHHAMMER: Plane analytical geometry, three hours; Differential equations with one independent variable. three hours; Seminar, one hour.—By Professor P. STAECKEL: Integral calculus with introduction to theory of differential equations, four hours; Abelian functions, four hours; Non-euclidean geometry, one hour; seminar, one hour.

UNIVERSITY OF KOENIGSBERG.—By Professor F. MEYER: Modern geometry of the triangle, one hour; Introduction to the theory of differential equations, four hours; Seminar, one hour.—By Professor A. SCHOENFLIES: Theory of functions, five hours; General introduction to higher mathematics, two hours; Seminar, two hours.—By Professor L. SAALSCHÜTZ: Differential calculus, two hours; Integral calculus, four hours, with exercises, one hour.—By Dr. E. MÜLLER: Descriptive geometry, four hours; Introduction to Grassmann's Ausdehnungslehre, two hours.

UNIVERSITY OF LEIPSIC.—Professor SCHEIBNER offers no courses this semester, and those of Professor Mayer will be announced later.—By Professor C. NEUMANN: Theory of potential and spherical harmonics, four hours; Seminar, two hours.—By Professor O. HÖLDER: Mechanics, four hours; Galois' theory of equations, two hours; Seminar, one hour. By Professor F. ENGEL: Infinitesimal calculus, four hours; Differential invariants and theory of groups, one hour; Seminar, one hour; Exercises in differential invariants and theory of groups, with Dr. Kowalewski, one hour.—By Dr. F. HAUSDORFF: Projective geometry treated synthetically, four hours; Non-euclidian geometry, two hours.—By Dr. G. KOWALEWSKI: Theory of functions, four hours, with exercises, one hour; Introduction to theory of transformation groups, two hours, with exercises, one hour.—By Dr. H. LIEBMANN: Analytical geometry of space, two hours, with exercies, one hour; Theory and application of determinants, two hours.

UNIVERSITY OF MARBURG.—By Professor F. SCHOTTKY: Integral calculus, four hours; Theory of numbers, four hours; Seminar, two hours.—By Professor E. HESS: Geometry of space considered analytically and synthetically, four hours; Theory of algebraic equations and determinants, four hours; Seminar, three hours.—By Dr. F. v. DALGWICK: Applications of infinitesimal calculus, three hours; Kinematics, one hour.

UNIVERSITY OF MUNICH.—By Professor G. BAUER: Plane analytical geometry; Seminar.—By Professor F. LINDEMANN: Differential calculus; Analytical mechanics; Line and sphere geometry; Seminar.—By Professor A. PRINGSHEIM: Elementary theory of infinite series, products, and continued fractions: Introduction to the theory of analytical functions.—By Dr. K. DÖHLEMANN: Descriptive geometry, with exercises; Synthetic geometry, with exercises.—By Dr. E. VON WEBER: Introduction to analysis; Differential equations.—By Dr. J. GÖTTLER: Algebra, with exercises.

UNIVERSITY OF STRASSBURG.—By Professor T. REYE: Geometry of space, modern methods, two hours; Mathematical theory of elasticity, two hours.—By Professor H. WEBER: Theory of numbers, four hours; Infinite series, two hours; Seminar in conjunction with Dr. Wellstein.—By Professor F. ROTH: Algebraic analysis and determinants, three hours; Analytical geometry of space, two hours; Ordinary differential equations, two hours.—By Professor A. KRAZER: Infinitesimal calculus, four hours; Plane analytical geometry, four hours: Seminar, two hours.—By Dr. E. TIMERDING: Theory of surfaces, two hours.—By Dr. J. WELLSTEIN: Graphical statics, two hours, with exercises, two hours.

UNIVERSITY OF TÜBINGEN.—By Professor A. VON BRILL: Introduction to higher mathematics, four hours; Theory of algebraic curves, three hours; Seminar, two hours.—By Professor H. STAHL: Higher analysis, II., four hours; Partial differential equations, three hours; Seminar, two hours.—By Professor L. MAURER: Descriptive geometry, II., one hour, with exercises, two hours; Calculus of variations, two hours, with exercises, one hour; Exercises in elementary geometry, one hour.

UNIVERSITY OF VIENNA.—By Professor R. v. ESCHERICH: Theory of functions five hours; Proseminar, one hour; Seminar, two hours.—By Professor L. GEGENBAUER: Elements of infinitesimal calculus, five hours, with exercises; Proseminar, one hour; Seminar, two hours.—By Professor F. MERTENS: Algebra, five hours; Proseminar, one hour; Seminar, one hour.—By Professor G. KOHN: Analytical geometry, four hours, with exercises, one hour; Differential geometry, II., one hour; The methods of descriptive geometry, one hour.—By Dr. V. SERSAWY: Lectures on the mathematics of insurance, two courses of three and four hours respectively.—By Dr. A. TAUBER: Mathematics of

insurance, four hours, with exercises, two hours.—By Dr. E. BLASCHKE: Mathematical statistics, 3 hours.—By Dr. R. D. v. STERNECK: Differential geometry, two hours; Kummer's theory of ideal numbers, one hour.—By Dr. K. CARDA: Theory of continuous groups, three hours.

OF the annual list of doctorates conferred by American universities during the academic year closing with June, 1901, *Science* records the following whose theses were in mathematics, the names of the universities conferring the degrees and the subjects of the dissertations being appended: GRACE ANDREWS, Columbia University, "The primitive double minimal surface of the seventh class and its conjugate"; C. W. McG. BLACK, Harvard University, "The parametric representation of the neighborhood of a singular point of an analytic function"; B. S. EASTON, University of Pennsylvania, "Substitutions and substitution groups"; W. FINDLAY, University of Chicago, "The Sylow subgroups of the symmetric group on k letters"; W. B. FITE, Cornell University, "On metabelian groups"; W. S. FRANKLIN, Cornell University, "Poynting's theorem"; A. S. GALE, Yale University, "On a particular class of algebraic minimum curves and surfaces"; C. N. HASKINS, Harvard University, "On the invariants of quadratic differential forms"; ARCHIBALD HENDERSON, University of South Carolina, "The cone of normals and an allied cone for central surfaces of the second degree"; L. I. HEWES, Yale University, "Some properties of path curves of continuous projective groups"; H. G. KEPPEL, Clark University, "The cubic 3-spread ruled with planes in 4-fold space"; H. W KUHN, Cornell University, "On imprimitive substitution groups"; CHARLOTTE E. PINGRA, University of Wisconsin, "On functions connected with special Riemann surfaces, in particular those for which $p = 3,4,5$"; I. E. RABINOVITCH, Johns Hopkins University, "The foundation of the euclidean geometry, as viewed from the standpoint of kinematics"; F. E. ROSS, University of California, "Differential equations belonging to a ternary linearoid group"; E. B. SKINNER, University of Chicago, "On ternary monomial substitution groups of finite order with determinant ± 1"; ROXANA H. VIVIAN, University of Pennsylvania, "The poles of a right line with respect to a curve of order n"; J. N. VAN DER VRIES, Clark University, "On the multiple points of twisted curves"; E. B. WILSON, Yale University, "The decomposition of the general collineation in space into three skew reflections";

RUTH G. WOOD, Yale University, "Non-euclidean displacements and symmetry transformations."

The list includes twenty names; the corresponding number in the year preceding was eleven.

THREE fasciculi of the Encyclopædia of the mathematical sciences have thus far appeared during the present calendar year. These are: Volume I, No. 6; Volume IV_2, No. 1; and Volume IV_1, No. 1. In press are: Volume I, No. 7 (concluding number); Volume II_1, No. 5; Volume II_2, No. 1; and Volume IV_1, No. 2. The *Archiv der Mathematik und Physik* has opened a special department for corrections, additions, and suggestions relative to the Encyclopædia. A considerable number of notes thus collected have been reprinted and inserted at the beginning of Volume IV_1, No. 1.

THE Yale bicentennial series, about to be issued by Charles Scribner's Sons, New York, includes two volumes representing mathematics: Elementary principles in statistical mechanics, by Professor J. WILLARD GIBBS; Vector analysis, a text book for the use of students of mathematics and physics, by Dr. E. B. WILSON.

MESSRS. LONGMANS AND CO. announce a text-book on "Higher mathematics for students of chemistry and physics" by Mr. J. W. MELLOR.

THE following are recent catalogues of new and second-hand works in the mathematical sciences: No. 205, S. Calvary and Co., Berlin; No. 28, Kirchoff and Wigand, Leipsic; No. 220, Albert Raustein, Zürich; No. 183, Mayer and Müller, Berlin; No. 77, Max Weg, Leipsic.

M. HERMANN, of Paris, is preparing a catalogue which is expected to appear in November.

THE firm of Gustav Fock, of Leipsic, has recently issued a catalogue containing about forty-five hundred titles of works in mathematics and physics drawn chiefly from the libraries of the late Professors OSCAR SCHLÖMILCH and ELWIN BRUNO CHRISTOFFEL.

ARRANGEMENTS were made by the Société des amis des sciences physiques et mathématiques of Poltava, Russia, to celebrate the centenary of the birth of MICHEL OSTROGRADSKY at Poltava on September 12–15.

PROFESSOR M. CANTOR, of the University of Heidelberg, recently celebrated the fifty year jubilee of his doctor's degree which he obtained at twenty-two years.

PROFESSOR J. J. THOMSON and Dr. J. LARMOR, of Cambridge University were among the recipients of the honorary degree of doctor of laws of the University of Glasgow conferred at the recent celebration of the ninth jubilee of that institution.

PROFESSOR W. W. JOHNSON, of the U. S. Naval Academy, has written a treatise on theoretical mechanics, which has just appeared from the press of John Wiley and Sons of New York.

A MEDALLION of the late Professor Charles Hermite will be placed in the Court of Honor of the Sorbonne.

PROFESSOR E. PICARD, of the Sorbonne, has been elected to foreign membership in the Reale Accademia dei Lincei of Rome.

PROFESSOR M. NÖTHER, of Erlangen, has been elected corresponding member of the Lombardy academy of sciences of Milan.

PROFESSOR F. ENGEL, of Leipsic, has received the gold Lobachevsky medal for his report (Gutachten) on Killing's work in connection with the award of the Lobachevsky prize by the University of Kazan (see BULLETIN, volume 7, p. 191).

DR. L. v. PRANTL, of Nuremberg, has been appointed professor of technical mathematics in the University of Jena.

PROFESSOR A. MAYER, of Leipsic, will not lecture during the coming winter semester, and will probably retire permanently.

DR. M. von RUDZKI has been promoted to a full professorship of mathematical geodesy at the University of Cracow.

DR. KARL CARDA has qualified as docent in mathematics at the University of Vienna, Dr. EDMUND LANDAU at the University of Berlin, and Dr. E. WEINHOLDT at the University of Kiel.

M. A. MANNHEIM, of the École Polytechnique of Paris, has retired from the active duties of his professorship after a tenure of thirty-seven years. His successor is M. HAAG.

MR. FREDERICK H. SEARES, recently instructor in astronomy at the University of California, has been appointed to the professorship in astronomy at the University of Missouri.

DR. J. M. PAGE has been promoted to a full professorship of mathematics at the University of Virginia.

DR. C. N. LITTLE, recently of Stanford University, has received the appointment of professor of civil engineering in the University of Idaho.

PROFESSOR T. C. ESTY, of Amherst College, has been appointed to the professorship of mathematics at the University of Rochester, succeeding Professor A. L. BAKER who has accepted a position in the Manual Training High School, Brooklyn, N. Y.

MR. PAUL ARNOLD has been appointed professor of mathematics in the University of Southern California.

DR. D. A. MURRAY, of Cornell University, has been appointed professor of mathematics in Dalhousie College, Halifax, Nova Scotia.

AT Ohio State University Professor J. E. BOYD has been promoted to an associate professorship of mathematics and Mr. D. E. RASOR has been appointed instructor in mathematics.

DR. H. F. STECKER has been appointed to an instructorship in mathematics at Cornell University.

DR. FREDERICK H. SAFFORD has resigned from the mathmatical staff of the University of Cincinnati.

DR. E. C. LUNN, of the University of Chicago, has been appointed instructor in mathematics and astronomy at Wesleyan University.

MR. S. W. REAVES has been appointed to an assistant professorship of mathematics in Clemson College, S. C.

MR. H. C. MORENO has been appointed instructor in mathematics in Stanford University.

MR. A. B. PIERCE, of the University of California, has received leave of absence and will spend the present academic year abroad.

DR. C. W. M. BLACK has been appointed instructor in mathematics in the University of Oregon.

PROFESSOR E. M. WOOD, of Baker University, Baldwin, Kansas, has been appointed professor of mathematics and astronomy at Albion College, Michigan, to succeed Professor HENRY BENNER, who was drowned at Lake Orion, August 14.

PAGES 39-52, SHOULD READ OCTOBER.

MR. CHARLES A. SCHOTT, chief of the computing division of the U. S. Coast and Geodetic Survey from 1855 to 1899, died at Washington, July 31, 1901.

THE deaths are announced of W. SCHUR, professor of astronomy of the University of Göttingen, aged 55 years; of Admiral DE JONQUIÈRES, of Paris, well known for his researches in geometry; of Mr. J. HAMBLIN SMITH, the author of numerous successful text-books in elementary mathematics, at the age of seventy-four years; of Professor P. HELMLING, of the University of Dorpat, at the age of 84.

PROFESSOR PETER GUTHRIE TAIT died at Edinburgh on July 4. He was born at Dalkeith in 1831 and was a student at Edinburgh University. Later he attended Cambridge University where he was senior wrangler, first Smith's prizeman and fellow of Peterhouse. In 1854 he became professor of mathematics at Queen's College, Belfast, whence he removed in 1860 to the chair of natural philosophy in the University of Edinburgh. He was the author of a long series of publications both technical and popular in character. His scientific papers were collected and published in 1898.

NEW PUBLICATIONS.

I. HIGHER MATHEMATICS.

ASCOLI (G.). See BRIOSCHI (F.).

BAUSCHINGER (J.). See ENCYKLOPÄDIE.

BELTRAMI (E.). See BRIOSCHI (F.).

BIANCHI (L.). Lezioni sulla teoria delle funzioni di variabile complessa e delle funzioni ellittiche. Pisa, Spoerri, 1901. 8vo. 608 pp. Fr. 20.00

BOHLMANN (G.). See ENCYKLOPÄDIE.

BOREL (E.). See CAUCHY (A.).

BORGMEYER (J.). Diskussion einer cyklischen Fläche dritter Ordnung. (Progr.) Duderstadt, 1901. 8vo. 29 pp.

BORTKIEWICZ (L. VON). See ENCYKLOPÄDIE.

BRIOSCHI (F.). Opere matematiche, pubblicate per cura del comitato per le onoranze a F. Brioschi (G. Ascoli, E. Beltrami, G. Colombo, L. Cremona, G. Negri, G. Schiaparelli). (In 4 volumi.) Vol. I. Milano, Hoepli, 1901. Folio, 12 + 416 pp., portrait. Fr. 25.00

BUHL (A.). Sur les équations différentielles simultanées et la forme aux dérivées partielles adjointe. (Thèse.) Paris, Naud, 1901. 8vo. 64 pp.

BURNSIDE (W. S.) and PANTON (A. W.). The theory of equations, with an introduction to the theory of binary algebraic forms. Vol. II. New York, Longmans, 1901. 8vo. 11 + 292 pp. Cloth. (Dublin University Press series.) $2.75

CANTOR (M.). Vorlesungen über Geschichte der Mathematik. Vol. III, Abt. 3, Abschnitt XVIII: 1725–1758. 2te Auflage. Leipzig, Teubner, 1901. 8vo. 10 pp. and pp. 493–923. M. 12.40.

CASALI (M.). See RASCHI (L.).

CAUCHY (A.). Oeuvres complètes, publiées sous la direction scientifique de l'Académie des sciences et sous les auspices du Ministère de l'instruction publique, avec le concours de MM. Valson, Collet et Borel. (En 27 volumes.) Première série: Mémoires, notes et articles extraits des recueils de l'Académie des sciences (12 volumes). Vol. XII. Paris, Gauthier-Villars, 1900. 4to. 504 pp. Fr. 25.00

CESÀRO (E.). Vorlesungen über natürliche Geometrie. Autorisierte deutsche Ausgabe von G. Kowalewski. Leipzig, Teubner, 1901. 8vo. 8 + 341 pp. Cloth. M. 12.00

COLLET. See CAUCHY (A.).

COLOMBO (G.), CREMONA (L.). See BRIOSCHI (F.).

CZUBER (E.). See ENCYKLOPÄDIE.

DEL RE (A.). Sulla struttura geometrica dello spazio in relazione al modo di percepire i fatti naturali. Discorso pronunziato in occasione della solenne inaugurazione degli studi presso la r. università di Modena il di 16 novembre 1896. 3a edizione, notevolmente arricchita di note. Napoli, Alvano, 1901. 8vo. 47 pp.

DICKSTEIN (S.). See PICARD (E.).

ENCYKLOPÄDIE der mathematischen Wissenschaften mit Einschluss ihrer Anwendungen. Herausgegeben im Auftrage der Akademien der Wissenschaften zu München und Wien und der Gesellschaft der Wissenschaften zu Göttingen, sowie unter Mitwirkung zahlreicher Fachgenossen. (In 7 Bänden.) Vol. I: Arithmetik und Algebra, redigiert von W. F. Meyer. Heft 6: H. Weber, Komplexe Multiplikation; E. Czuber, Wahrscheinlichkeitsrechnung; J. Bauschinger, Ausgleichungsrechnung (Methode der kleinsten Quadrate, Fehlertheorie); J. Bauschinger, Interpolation; L. von Bortkiewicz, Anwendungen der Wahrscheinlichkeitsrechnung auf Statistik; G. Bohlmann, Lebensversicherungs-Mathematik; D. Seliwanoff, Differenzenrechnung; R. Mehmke, Numerisches Rechnen. Leipzig, Teubner, 1901. 8vo. Pp. 721–992. M. 7.20

ESTANAVE (E.). Sur une série servant à définir approximativement le nombre π, rapport de la circonférence au diamètre. Paris, Croville-Morant, 1901. 8vo. 16 pp.

FRICKE (R.) und KLEIN (F.). Vorlesungen über die Theorie der automorphen Functionen. Vol. II: Die functionentheoretischen Ausführungen und die Anwendungen. Lieferung 1: Engere Theorie der automorphen Functionen. Leipzig, Teubner, 1901. 8vo. 6 + 282 pp. M. 10.00

GOLDBERG (A.). Die jüdischen Mathematiker und die jüdischen anonymen mathematischen Schriften, alphabetisch geordnet mit Angabe ihrer Zeit, zugleich ein Index zu M. Steinschneider's Mathematik bei den Juden. Frankfurt a. M., Kauffmann, 1901. 8vo. 12 pp. M. 1.00

GORDAN (P.). Die Hessesche und die Cayleysche Kurve. Leipzig, Deichert, 1901. 8vo. 14 pp. (Aus "Festschrift der Universität Erlangen für Prinzregent Luitpold.") M. 0.60

GRÜNBAUM (H.). Lehr- und Uebungsbuch der Differentialrechnung, enthaltend über 750 Uebungsaufgaben, für mittlere technische Lehranstalten, Realgymnasien, Oberrealschulen u. s. w., sowie zum Selbststudium. Würzburg, Frank, 1901. 12mo. 6 + 137 pp. M. 2.80

GUTZMER (A.). See JAHRESBERICHT.

HAENTZSCHEL (E.). Elementare Herleitung der Newtonschen Reihen für Sinus and Cosinus und die Normierung der Vorzeichen bei der Definition der trigonometrischen Funktionen. (Progr.) Berlin, Gaertner, 1901. 4to. 22 pp. M. 1.00

HEDRICK (E. R.). Ueber den analytischen Character der Lösungen der Differentialgleichungen. (Diss., Göttingen, 1901.) Göttingen, Dieterich, 1901. 8vo. 78 pp.

HENSEL (K.). See KRONECKER (L.).

JAHRESBERICHT der deutschen Mathematiker-Vereinigung. Vol. 5: Enthaltend die Chronik der Vereinigung für das Jahr 1896, die auf der Versammlung in Frankfurt a. M. gehaltenen Vorträge, sowie: Die Entwickelung der synthetischen Geometrie. Teil I: Von Monge bis auf Staudt (1847). Bericht von Ernst Kötter in Aachen. Herausgegeben von A. Wangerin und A. Gutzmer. 2te (Schluss-) Lieferung. Leipzig, Teubner, 1901. 8vo. 28 pp. and pp. 129–486. M. 14.40

IVEY (J. N.). Ueber die Periodicitätsmoduln der Abelschen Integrale erster Gattung. (Diss.) Berlin, 1901. 8vo. 52 pp. M. 1.80

KAPTEYN (W.). De meer-dimensioneele meetkunde. Utrecht, 1901. 8vo. 4 + 29 pp. M. 1.00

KASNER (E.). The invariant theory of the inversion group; geometry upon a quadric surface. 1900. 4to. (*Transactions of the American Mathematical Society*, 1, pp. 430–498.)

KLEIN (F.). See FRICKE (R.).

KÖTTER (E.). Die Entwickelung der synthetischen Geometrie von Monge bis auf Staudt (1847). Erster Band eines Berichtes, erstattet der deutschen Mathematiker-Vereinigung. Leipzig, Teubner, 1901. 8vo. 28 + 486 pp. M. 18.80

—— See JAHRESBERICHT.

KOWALEWSKI (G.). See CESÀRO (E.).

KRONECKER (L.). Vorlesungen über Mathematik, herausgegeben unter Mitwirkung einer von der k. preussischen Akademie der Wissenschaften eingesetzten Kommission. (In 2 Teilen.) Teil 2: Vorlesungen über allgemeine Arithmetik, bearbeitet und herausgegeben von K. Hensel. Erster Abschnitt: Vorlesungen über Zahlentheorie. Vol. I. Leipzig, Teubner, 1901. 8vo. 13 + 509 pp. M. 16.00

KRUMME (W.). Der Unterricht in der analytischen Geometrie. Für Lehrer und zum Selbstunterricht. Berlin, Salle, 1901. 8vo. M. 6.50

LIEBE (A.). Ueber die Analogie der aus der Entwickelung von $(1-2ax+a^2)^{-\lambda}$ entspringenden Funktionen mit den Kegelfunktionen. II. (Progr.) Borna, 1901. 4to. 24 pp.

LOOMIS (E.). The elements of analytical geometry. Revised edition. New York, American Book Company, [1900]. 12mo. 261 pp. Shp. $1.00

LORENTZ (H.). The foundations of higher mathematics: the elements of analytic geometry, the differential and integral calculus, and their applications in the natural sciences. Freely translated and accompanied by an historical sketch of the development of mathematical analysis by V. P. Sheromedevsky. Vol. II. Moscow, 1901. 8vo. 620 pp. (Russian.) M. 7.50

MEHMKE (R.), MEYER (W. F.). See ENCYKLOPÄDIE.

MOJARES (U.). See RASCHI (L.).

MÜLLER (F.). Mathematisches Vokabularium französisch-deutsch und deutsch-französisch, enthaltend die Kunstausdrücke aus der reinen und angewandten Mathematik. 2te Hälfte. Leipzig, Teubner, 1901. 8vo. Pp. ix–xiv and 133–315. M. 11.00

NEGRI (G.). See BRIOSCHI (F.).

NERNST (W.) und SCHOENFLIES (A.). Einführung in die mathematische Behandlung der Naturwissenschaften. Kurzgefasstes Lehrbuch der Differential- und Integralrechnung mit besonderer Berücksichtigung der Chemie. 3te Auflage. München, Wolff, 1901. 8vo. 12 + 340 pp. M. 10.00

NOETHER (M.). Zur Erinnerung an Karl Georg Christian von Staudt. Leipzig, Deichert, 1901. 8vo. 24 pp. (Aus "Festschrift der Universität Erlangen für Prinzregent Luitpold.") M. 0.80

OSGOOD (W. F.). Note über analytische Functionen mehrerer Veränderlichen. 1900. 8vo. (*Mathematische Annalen* 52, pp. 462–464.)

——. Zweite Note über analytische Functionen mehrerer Veränderlichen. 1901. 8vo. (*Mathematische Annalen* 53, pp. 461–464.)

——. On a fundamental property of a minimum in the calculus of variations and the proof of a theorem of Weierstrass's. 1901. 4to. (*Transactions of the American Mathematical Society* 2, pp. 273–295.)

PANTON (A. W.). See BURNSIDE (W. S.).

PICARD (E.). L'œuvre scientifique de Charles Hermite. Leçon faite à la Faculté des sciences, le samedi 2 mars 1901. 4to. 26 pp., portrait. (*Annales de l'Ecole normale supérieure* (3) 18.)

——. O rozwoju niektórych teoryj zasadniczych analizy matematycznéj w wieku XIX. Przełożył S. Dickstein. Warszawa, 1900. 8vo. 59 pp. M. 2.00

RADELFINGER (F. G.). Progress of pure mathematics in 1900. Washington, D. C., 1901. 8vo. (*Bulletin of the Philosophical Society of Washington*, March, 1901, pp. 157–165.)

RASCHI (L.). Algebra complementare. Appunti compilati da M. Casali e U. Mojares sulle lezioni [tenute nella] r. università di Parma, 1900–1901. . Parma, Zafferri, 1900–1901. 8vo. Disp. 1–10, pp. 1–78.

——. Geometria analitica. Appunti compilati da U. Mojares e M. Casali sulle lezioni [tenute nella] r. università di Parma. Parma, Zafferri, 1900–1901. 8vo. Disp. 1–11, pp. 1–86.

RIEMANN (B.). See WEBER (H.).

SALA (L.). La proporzionalità nel calcolo cosi elementare come differenziale ed integrale. Milano, Tipografia degli ingegneri, 1901. 8vo. 17 pp.

SCHAEWEN (P. VON). Die Binomialkoefficienten in Verbindung mit figurierten Zahlen und arithmetischen Reihen höherer Ordnung. Neuer Abdruck. Glogau, Flemming, 1901. 8vo. 30 pp. M. 1.20

SCHIAPARELLI (G.). See BRIOSCHI (F.).

SCHOENFLIES (A.). See NERNST (W.).

SCHOLIM (P.). Einige Eigenschaften einer ebenen Kurve vierter Ordnung und einer ihr entsprechenden Raumkurve. (Progr.) Königshütte, O.-S., 1901. 4to. 9 pp.

SCHOUTEN (G.). Analytische meetkunde von het platte vlak (Kegelsneden). 2te druk. Delft, 1901. 8vo. 146 pp. M. 4.50

SELIWANOFF (D.). See ENCYKLOPÄDIE.

SHEROMEDEVSKY (V. P.). See LORENTZ (H.).

VALSON. See CAUCHY (A.).

VAN VLECK (E. B.). On the convergence of continued fractions. 1901. 4to. (*Transactions of the American Mathematical Society* 2, pp. 215–233.)

VIVIAN (R. H.). The poles of a right line with respect to a curve of order n. (Thesis, University of Pennsylvania.) Philadelphia, 1901. 8vo. 32 pp.

WANGERIN (A.). See JAHRESBERICHT.

WEBER (H.). Die partiellen Differential-Gleichungen der mathematischen Physik. Nach Riemann's Vorlesungen in vierter Auflage neu bearbeitet. (In 2 Bänden.) Vol. II. Braunschweig, Vieweg, 1901. 8vo. 12 + 527 pp. M. 10.00

WEBER (H,). See ENCYKLOPÄDIE.

WILCZYNSKI (E. J.). Transformation of systems of linear differential equations. 1901. 4to. (*American Journal of Mathematics* 23, pp. 29–36.)

II. ELEMENTARY MATHEMATICS.

ADAMOVICH (S.). Theory of the resolution of algebraic expressions into factors, with examples. St. Petersburg, 1901. 8vo. 47 pp. (Russian.) M. 2.00

ALBUQUERQUE (J. D'A.). Trigonometria plana de conformidade com o programma official do ensino secundario e com 151 exercicios. Porto, Imprensa portugueza, 1901. 8vo. 202 pp.

ARNALDI (M.). Prime nozioni di geometria intuitiva, ad uso delle scuole complementari. Parma, Battei, 1901. 16mo. 160 pp. Fr. 1.50

BAUR (L.). Untersuchungen über die verschiedenen Wurzeln einer algebraischen Gleichung. (Progr.) Heppenheim, 1901. 8vo. 32 pp.

BERNARDI (L.). Elementi di trigonometria piana. Udine, Tosolini, 1901. 16mo. 132 pp.

BERTRAND (J.). Traité d'arithmétique. 12e édition. Paris, Hachette, 1901. 8vo. 344 pp. Fr. 4.00

——. Theoretical arithmetic, presented mainly according to Bertrand by N. Bilibin. 5th edition. St. Petersburg, 1901. 8vo. 292 pp. (Russian.) M. 4.00

BEURIGER (J.). Zur Auflösung der biquadratischen Gleichungen. (Progr.) Bonn, 1901. 4to. 20 pp.

BILIBIN (N.) See BERTRAND (J.).

BRAUER (E. A.). Springende Logarithmen. Abgekürzte fünfstellige Logarithmentafel mit zunehmenden Grundzahlstufen, zum Gebrauch für technische Rechnungen bearbeitet. Karlsruhe, 1901. Boards. M. 0.90

BRIOT (C.). Algebra elemental y superior. Traducida, anotada y completada por C. Sebastian y B. Portuondo. Nueva edición. Madrid, 1900. 4to. 732 pp., 2 plates. M. 14.00

BROOKSMITH (E. J.). Woolwich mathematical papers for admission to the royal military academy, for the years 1891 to 1900. London, 1901. 8vo. Cloth. 6s.

BRYANT (S.). See SMITH (C.).

BÜTZBERGER (F.). Lehrbuch der ebenen Trigonometrie, mit vielen angewandten Aufgaben für Gymnasien und technische Mittelschulen. 2te, umgearbeitete und erweiterte Auflage. Zürich, Füssli, 1901. 8vo. 8 + 62 pp. Boards. M. 2.00

CHAILAN (E.). Résumé d'algèbre élémentaire, à l'usage des classes de lettres et des candidats à la première partie du baccalauréat de l'enseignement secondaire classique. 4e édition. Paris, Poussielgue, 1901. 18mo. 73 pp. (Alliance des maisons d'éducation chrétienne.)

DEAKIN (R.). Tutorial algebra. Part I: Elementary course. London, Clive, 1901. 12mo. 452 pp. Cloth. (University tutorial series.) 3s. 6d.

DENNERT (E.). Mathematisches Formelbuch. Godesberg, Schugt, 1901. 8vo. 36 pp. M. 0.75

EBNETER (K.). Geometrie für Sekundarschulen. Heft I. 4te Auflage des "Leitfadens." St. Gallen, Fehr, 1901. 8vo. 96 pp. Boards. M. 1.20

EDMONDSON (T. W.). Deductions in Euclid. London, Clive, 1901. 12mo. 210 pp. Cloth. (University tutorial series.) 2s. 6d.

EUCLID. See EDMONDSON (T. W.), LARMOR (A.), SMITH (C.).

FENKNER (H.). Arithmetische Aufgaben, unter besonderer Berücksichtigung von Anwendungen aus dem Gebiete der Geometrie, Physik und Chemie. Ausgabe A, vornehmlich für den Gebrauch in Gymnasien, Realgymmasien und Oberrealschulen. Teil I: Pensum der Untertertia, Obertertia und Untersekunda. 4te Auflage. Berlin, Salle, 1901. 8vo. 8 + 256 pp. M. 2.20

FLOR (O.). Lösung des Problems: die Quadratur des Kreises; Berichtigung der Zahl π. 2te Ausgabe. Riga, Jonck and Poliewsky, 1901. 8vo. 8 pp. M. 2.00

FRIOCOURT (G.). Tables de logarithmes à six décimales pour les nombres et les lignes trigonométriques, et tables de navigation. Nouvelle édition, révisée. Paris, 1901. 8vo. Fr. 10.00

GELIN (E.). Précis d'arithmétique élémentaire, à l'usage des écoles moyennes, des cours professionnels, des écoles normales et des classes d'humanités. Ouvrage adopté par le conseil de perfectionnement de l'enseignement moyen pour les écoles moyennes, les collèges et les athénées. 4e édition, rédigée d'après un plan simplifié, et contenant les solutions raisonnées de 130 problèmes-types. Namur, Wesmael-Charlier, 1900. 8vo. 272 pp. Fr. 3.00

——. Su di un sistema di equazioni del primo grado. 8vo. 8 pp. (*Le matematiche pure ed applicate*, I, 1901.)

GREMIGNI (M.). Elementi di geometria ad uso delle scuole tecniche e professionali. Planimetria, libro 3–4. Firenze, Bemporad, 1901. 16mo. 7 + 120 pp. (Biblioteca scolastica.) Fr. 0.80

HADAMARD (J.). Leçons de géométrie élémentaire. Partie II: Géométrie dans l'espace. Paris, Colin, 1901. 8vo. 21 + 582 pp. (Cours complet de mathématiques élémentaires.) Fr. 10.00

HALLER VON HALLERSTEIN (F.). Lehrbuch der Elementarmathematik. Nach dem Lehrplane für das k. preussische Kadettenkorps bearbeitet von B. Hülsen. Teil 2: Pensum der Obertertia. 4te Auflage. Berlin, Nauck, 1901. 8vo. 8 + 192 pp. Hf. leather. M. 3.40

HAMELIN (G.). See NAUD (L.).

HESSENBERG (G.). Ebene und sphärische Trigonometrie. Neudruck. Leipzig, Göschen, 1901. 12mo. 165 pp. Cloth. (Sammlung Göschen, No. 99.) M. 0.80

HILL (G. A.). See WENTWORTH (G. A.).

HOLGATE (T. F.). Elementary geometry, plane and solid. For use in high schools and academies. New York, Macmillan, 1901. 12mo. 11 + 440 pp. Hf. leather. $1.10

HÜLSEN (B.). See HALLER VON HALLERSTEIN (F.).

JENTZEN (E.). Elemente der Trigonometrie zum praktischen Gebrauch für Unterrichtazwecke an mittleren technischen Lehranstalten. 3te Auflage. Dresden, Kühtmann, 1901. 8vo. 63 pp. M. 1.60

KINKELIN (H.). See RIEM (J.).

KÖLTZSCH (A.). Grundzüge der Raumlehre. Ein Lern- und Uebungsbuch, zum Gebrauche in Volksschulen, Fortbildungsschulen, Präparantenanstalten und Mittelschulen. Heft II. Mit 80 Textfiguren und mehr als 300 verschiedenen Uebungsaufgaben. 2te Auflage. Leipzig, Merseburger, 1901. 8vo. β + 74 pp. M. 0.60

LARMOR (A.). Geometrical exercises from Nixon's "Euclid revised," with solutions. Oxford, Clarendon Press, 1901. 12mo. 8 + 172 pp. Cloth. $0.90

LONDON MATRICULATION MATHEMATICS: Guide to mathematics of University matriculation examinations. 8th edition. London, Clive, 1901. 12mo. 108 pp. (University tutorial series.) 2s.

M. (F. G.). Exercices d'algèbre. 5e édition. Paris, Poussielgue, 1901. 8vo. 24 + 955 pp. (Cours de mathématiques élémentaires.)

MEISTER (F.). Grundriss der Geometrie. Zum Gebrauche an Sekundarschulen bearbeitet. Zürich, Schulthess, 1901. 8vo. 78 pp. Boards. M. 0.80

——. Schlüssel zu den Aufgaben. 14 pp. M. 0.60

MOČNIK (F. VON). Lehrbuch der Arithmetik und Algebra nebst einer Aufgabensammlung für die oberen Klassen der Mittelschulen, bearbeitet von A. Neumann. Ausgabe für Gymnasien. 26ste Auflage. Wien und Prag, Tempsky, 1901. 8vo. 312 pp. Cloth. M. 3.70

MOST (R.). Der mathematische Unterrichtsstoff und das mathematische Bildungsgebiet in den oberen Klassen des Realgymnasiums und der Oberrealschule. (Progr., Coblenz.) Leipzig, Fock, 1901. 8vo. 8 + 200 + 26 pp., 22 plates. M. 4.00

NAUD (L.) et HAMELIN (G.). Cours résumé de mathématiques (arithmétique, système métrique, géométrie et algèbre), à l'usage des candidats aux examens de l'administration des postes et des télégraphes. Paris, bureaux du *Courrier des examens*, [1901]. 8vo. 208 pp. (Bibliothèque du *Courrier des examens des postes, des télégraphes et des téléphones.*) Fr. 2.50

NEUMANN (A.). See MOČNIK (F. VON).

NICHOLS (W. F.). Arithmetical problems. Boston, Thompson, Brown & Co., 1901. 12mo. 8 + 102 pp. Cloth. $0.30

NIEHUS (P.). Die Elemente der Arithmetik und der Algebra. Für Baugewerkschulen, Maschinenbauschulen, Handwerkerschulen und sonstige Berufsanstalten. Magdeburg, Friese, 1901. 8vo. 112 pp. Boards. M. 2.00

NIXON. See LARMOR (A.).

ORTU CARBONI (S.). Esercizi di geometria elementare, oltre 1350, con razionali avviamenti alle soluzioni. Livorno, Giusti, 1901. 16mo. 8 + 170 pp. (Biblioteca degli studenti; riassunti per tutte le materie d'esame nei licei, ginnasi, istituti tecnici, ecc., Vol. 54–55.) Fr. 1.00

OTTE (A.). Aufgaben über das gleichschenklige Dreieck in algebraischer Behandlung. (Progr.) Delitzsch, 1901. 4to. 14 pp.

PORTUONDO (B.). See BRIOT (C.).

PUCCINI (E.). Il concetto dell' infinitesimo nello studio della matematica elementare. Lettera aperta a Francesco Torraca. Pistoia, Flori, 1901. 8vo. 22 pp., 2 plates. Fr. 0.50

RIEM (J.). Rechentabellen für Multiplikation und Division. Mit Vorwort von H. Kinkelin. 2te Auflage. München, 1901. 4to. 101 pp. M. 6.00

RÜEFLI (J.). Kleines Lehrbuch der ebenen Geometrie nebst einer Sammlung von Uebungsaufgaben. Zum Gebrauche an Mittelschulen bearbeitet. 5te Auflage. Bern, Schmid & Francke, 1901. 8vo. 4 + 84 pp. Boards. M. 0.80

RUPERT (W. W.) Famous geometrical theorems and problems, with their history. (In 4 parts.) Part IV. Boston, Heath, 1901. 16mo. Pp. 76–107.

SALOMON (MME A.). Leçons d' algèbre, à l'usage de l'enseignement secondaire des jeunes filles (classes de quatrième et cinquième années) et des aspirants au brevet supérieur. Paris, Nony, 1901. 18mo. 188 pp.

SAVIN (M. S.). Solutions of the problems of the quadrature of the circle and of the trisection of the angle; construction of a square of equal magnitude of that of a circle and exact division of an angle into three equal parts, by means of propositions of elementary geometry. Orenburg, 1901. 8vo. 26 pp., 1 plate. (Russian.) M. 2.50

SCHERER. Zweistellige Multiplicationstafel zum Gebrauch im Büreau, Comtoir u. s. w. Cassel, Scherer, 1901. 4 sheets folio. M. 2.00

SCHMEHL (C.). Die Algebra und algebraische Analysis, mit Einschluss der elementaren Theorie der Determinanten in den oberen Klassen von höheren Lehranstalten, insbesondere der Realgymnasien und Oberrealschulen. Für den Schulgebrauch und Selbstunterricht bearbeitet. Giessen, Roth, 1901. 8vo. 8 + 286 pp. M. 2.50

SCHOTSCH (F.). Mathematische Aufgaben für die Maturitätsprüfung. Mediasch, 1901. 4to. 38 pp.

SCOTTI (G.). Elementi di geometria, ad uso del ginnasio superiore, secondo gli ultimi programmi governativi. Torino, Tipografia Salesiana, 1901. 8vo. 128 pp. Fr. 1.50

——. Elementi di geometria intuitiva ad uso del ginnasio inferiore e dei corsi complementari, secondo gli ultimi programmi governativi. Torino, Tipografia Salesiana, 1901. 8vo. 139 pp. Fr. 1.00

SEBASTIAN (C.). See BRIOT (C.).

SMITH (C.) and BRYANT (S.). Euclid's Elements of geometry, books 1–4, 6, 11. Edition for schools. London, Macmillan, 1901. 12mo. 468 pp. Cloth. 4s. 6d.

——. Books 6 and 11. Edition for schools. London, Macmillan, 1901. 12mo. 177 pp. Cloth. 1s. 6d.

SPIEKER (T.). Lehrbuch der Stereometrie mit Uebungsaufgaben für höhere Lehranstalten. 3te Auflage, 4tes bis 6tes Tausend. Potsdam, Stein, 1901. 8vo. 4 + 119 pp. Cloth. M. 2.00

STRAKHOV (M. A.). Short text-book of geometry, with practical applications. St. Petersburg, 1901. 8vo. 144 pp. (Russian.) M. 2.50

VERONESE (G.). Nozioni elementari di geometria intuitiva pei ginnasi inferiori. Padova, Drucker, 1901.

WAGNER (H.). Die ebene Trigonometrie in rein geometrischer Behandlung. (Progr.) Hamburg, 1901. 4to. 20 pp., 1 plate.

WEHNER (H.). Leitfaden für den stereometrischen Unterricht an Realschulen. 2te Auflage. Leipzig, Teubner, 1901. 8vo. 5 + 65 pp. Boards. M. 1.00

WENTWORTH (G. A.) and HILL (G. A.). The first steps in geometry. Boston, Ginn, 1901. 12mo. 4 + 156 pp. Hf. leather. $0.65

WILK (E.). Der gegenwärtige Stand der Geometrie-Methodik. Dresden, 1901. 8vo. 3 + 81 pp. M. 1.30

ZWICKY (M.). Grundriss der Planimetrie und Stereometrie, nebst Uebungsaufgaben. Teil 2: Stereometrie. 2te Auflage. Bern, Schmid & Francke, 1901. 8vo. 5 + 67 pp. Boards. M. 1.20

III. APPLIED MATHEMATICS.

ABRAHAM (M.). See ENCYKLOPÄDIE.

BEAU (O.). Die Berechnung der Sonnen- und Mondfinsternisse. III: Die ausführliche Berechnung der Sonnenfinsternisse. (Progr.) Sorau, 1901. 4to. 14 pp., 1 plate.

BIGOURDAN (G.). Le système métrique des poids et mesures; son établissement, sa propagation graduelle; histoire des opérations qui ont servi à déterminer le mètre et le kilogramme. Paris, Gauthier-Villars, 1901. 8vo. 456 pp., 10 plates. Fr. 10.00

BISKE (F.). Versuch einer Anwendung hydrodynamischer Untersuchungen auf die Protuberanzen der Sonne. (Diss.) Berlin, 1901. 8vo. 37 pp. M. 1.60

BORCK (R.). Interferenzkurven eines Wellensystems, welches mit einer Phasenverzögerung an einer festen Wand reflektiert wird. (Progr.) Liegnitz, 1901. 8vo. 22 pp.

BREITHOF (F.). Stéréotomie; théorie et construction des arches biaises. Paris, Gauthier-Villars, 1901. 8vo. 68 pp., with a 4to atlas of 10 plates. Fr. 5.50

BREITHOF (N.) et BREITHOF (F.). Traité de géométrie descriptive. Partie I: Point, droite, plan. 4e édition, revue et augmentée. Paris, Gauthier-Villars, 1901. 8vo. 193 pp., with 4to atlas of 24 plates. Fr. 11.50

BRUNELLI (P. H.). Calcul et construction des coupoles métalliques réticulaires. Traduit de l'italien par D. Mathieu. Paris, Dunod, 1901. 8vo. 9 + 216 pp.

CARVALLO (E.). Traité de mécanique, à l'usage des élèves de mathématiques élémentaires, des aspirants au baccalauréat de l'enseignement classique (2e série) et au baccalauréat de l'enseignement moderne (3e série), et des candidats à l'Institut agronomique et à l'Ecole navale. 2e édition. Paris, Nony, 1901. 8vo. 4 + 162 pp.

CAVEGLIA (C.). Sulla teoria delle travi e dei lastroni di cemento armato caricati di pesi. Roma, Voghera, 1900. 8vo. 98 pp., 2 plates.

CLAUSSEN (E.). Die statische Berechnung der Fabrikschornsteine. Auf Grund amtlichen Materials allgemeinverständlich dargestellt und für den praktischen Gebrauch bearbeitet. 2te Auflage. Lüneburg, König, 1901. 8vo. 4 + 79 pp., 1 plate. M. 3.00

DINSE (P.). See GELCICH (E.).

ENCYKLOPÄDIE der mathematischen Wissenschaften mit Einschluss ihrer Anwendungen. Herausgegeben im Auftrage der Akademien der Wissenschaften zu München und Wien und der Gesellschaft der Wissenschaften zu Göttingen, sowie unter Mitwirkung zahlreicher Fachgenossen. (In 7 Bänden.) Vol IV: Mechanik, regidiert von F. Klein. (In 2 Teilen.) Teil 2. Heft 1: M. Abraham, Geometrische Grundbegriffe; A. E. H. Love, Hydrodynamik: physikalische Grundlegung; A. E. H. Love, Hydrodynamik: theoretische Ausführungen. Leipzig, Teubner, 1901. 8vo. 147 pp. M. 3.80

FÖPPL (A.). Vorlesungen über technische Mechanik. (In 4 Bänden.) Vol. IV: Dynamik. 2te Auflage. Leipzig, Teubner, 1901. 8vo. 15 + 506 pp. Cloth. M. 12.00

GEDICUS (F. W.). Kinetik. Beiträge zu einer einheitlichen mechanischen Grundanschauung. Wiesbaden, Bergmann, 1901. 8vo. 12 + 124 pp. M. 2.40

GELCICH (E.) und SAUTER (F.). Kartenkunde, geschichtlich dargestellt. 2te Auflage, besorgt von P. Dinse. 2ter Abdruck. Leipzig, Göschen, 1901. 12mo. 168 pp. Cloth. (Sammlung Göschen, No. 30.) M. 0.80

GRAWINKEL (C.). See THOMPSON (S. P.).

GRÜNEWALD (C.). Zur Mathieuschen Theorie der Traversalschwingungen elastischer Scheiben und ihrer Prüfung durch Barthélemy. (Progr.) Berlin, 1901. 4to. 24 pp.

HERRMANN (G.). See WEISBACH (J.).

HOPKINSON (J.). Original papers, edited by B. Hopkinson. (Cambridge University press series.) Vol. I: Technical papers, with a memoir. London, Macmillan, 1901. 8vo. 360 pp. Cloth. $3.50

——. Vol. II: Scientific papers. London, Macmillan, 1901. 8vo. 400 pp. Cloth. $3.50

JOHNSON (W. W.). Theoretical mechanics; an elementary treatise. New York, Wiley, 1901. 12mo. 15 + 434 pp. Cloth. $3.00

KECK (W.). Fragen über die wichtigsten Gegenstände aus dem Gebiete der Mechanik. 4te Auflage. Hannover, Helwing, 1901. 8vo. 16 pp. M. 0.50

KLEIN (F.). See ENCYKLOPÄDIE.

KOLL (O.). Die Theorie der Beobachtungsfehler und die Methode der kleinsten Quadrate mit ihrer Anwendung auf die Geodäsie und die Wassermessungen. 2te, vermehrte und verbesserte Auflage. Berlin, Springer, 1901. 8vo. 12 + 323 + 31 pp. M. 10.00

KORN (A.). Eine mechanische Theorie der Reibung in kontinuierlichen Massensystemen. Berlin, Dümmler, 1901. 8vo. 12 + 219 pp. M. 6.00

KRÜGER (J.). Ueber den Einfluss der Temperatur auf die Wärmeleitung von Gläsern. (Diss.) Jena, 1901. 8vo. 44 pp.

LEBON (E.). Traité de géométrie descriptive et de géométrie cotée. Vol. I. Paris, Delalain, 1901. 8vo. 6 + 280 pp. Fr. 5.00

LEITFADEN für den Unterricht in der Navigation. Auf Veranlassung der Inspektion des Bildungswesens der Marine bearbeitet. 3te Auflage, mit Anhang: Muster zu nautischen Rechnungen. Berlin, 1901. 8vo. M. 12.50

LORENTZ (H. A.). Zichtbare en onzichtbare bewegingen. Leiden, 1901. 8vo. 4 + 173 pp. Fr. 2.50

LOVE (A. E. H.). See ENCYKLOPÄDIE.

LUBELZA (J.). Apuntes para una mecánica fundamental sintética. Madrid, 1901. 8vo. 73 pp.

LYONS (T. A.). A treatise on electromagnetic phenomena and on the compass and its deviations aboard ship, mathematical, theoretical, and practical. Vol. I. New York, 1901. 8vo. 15 + 556 pp. $6.00

MACH (E.). Die Mechanik in ihrer Entwickelung historisch-kritisch dargestellt. 4te Auflage. Leipzig, Brockhaus, 1901. 8vo. 14 + 550 pp. (Internationale wissenschaftliche Bibliothek, Vol. 59.) M. 8.00

MARC (L.). Sammlung der Aufgaben aus der höheren Mathematik, technischen Mechanik und darstellenden Geometrie, welche bei der Vorprüfung für das Bauingenieur-, Architektur- und Maschinen-Ingenieurfach an der k. technischen Hochschule zu München in den Jahren 1885 bis 1901 gestellt worden sind. München, Ackermann, 1901. 8vo. 3 + 52 pp. M. 1.60

MATHIEU (D.). See BRUNELLI (P. H.).

MOORE (E. C. S.). New tables for the complete solution of Ganguillet and Kutter's formula for the flow of liquid in open channels, pipes, sewers, and conduits. New York, Van Nostrand, 1901. 8vo. Cloth. $5.00

MÜLLER-BRESLAU (H. F. B.). Die graphische Statik der Baukonstruktionen. 3te Auflage. Vol. I: Zusammensetzung und Zerlegung der Kräfte in der Ebene; Trägheitsmomente und Centrifugalmomente ebener Querschnitte; Spannungen in geraden Stäben; Theorie der statisch bestimmten Träger mit Ausschluss der Untersuchung der Formänderungen. Leipzig, Baumgärtner, 1901. 8vo. 8 + 554 pp., 7 plates. M. 18.00

OLSHAUSEN (G. R.). Ueber die Unipolarrotation. (Diss.) Berlin, 1901. 4to. 43 pp. M. 2.40

PERRIN (E.). Nouvelles tables destinées à abréger les calculs nautiques. 3e édition, révisée. Paris, 1901. 8vo. Fr. 4.00

PLATNER (G.). Die Mechanik der Atome. Berlin, Krayn, 1901. 8vo. 4 + 97 pp. M. 2.50

PLESSEN (K. VON). Ueber den Einfluss suspendierter Teilchen auf den Auftrieb einer Flüssigkeit. (Diss.) Greifswald, 1901. 8vo. 40 pp.

SAUTER (F.). See GELCICH (E.).

SCHILLER (N.). Der Begriff des thermischen Verkehrs als Grundlage des zweiten thermodynamischen Hauptsatzes. Leipzig, Barth, 1901. 8vo. (*Annalen der Physik* (4) 5, pp. 313–325.)

——. Zur Thermodynamik gesättigter Lösungen. Leipzig, Barth, 1901. 8vo. (*Annalen der Physik* (4) 5, pp. 326–348.)

SCHÖNEMANN. Ueber die Ermittelung von Entfernungen und Höhen durch perspektivische Beziehungen. (Progr.) Soest, 1901. 4to. 32 pp., 2 plates.

SCHRAMM (W.). Ueber die Verteilung des Lichts in der Atmosphäre. (Diss.) Kiel, 1901. 8vo. 51 pp., 2 plates.

SCHUBERT (H.). Theorie des Schlickschen Massenvergleichs bei mehrkurbeligen Dampfmaschinen. Leipzig, 1901. 8vo. 132 pp. M. 12.00

SCHULZE (P.). Ueber asymmetrische Schwingungen um eine Lage stabilen Gleichgewichts. (Diss.) Greifswald, 1901. 8vo. 97 pp.

SCHWARZ (A.). Die Bilanzrechnung bei Pensionsinstituten. Wien, 1901. 8vo. 15 pp. M. 1.00

SERVUS (H.). Die Störungen der Atmosphäre und des Erdinnern durch Sonne und Mond. Neue Grundlagen der Meteorologie. Teil 2. (Progr.) Berlin, Gaertner, 1901. 4to. 18 pp. M. 1.00

SPANGLER (H. W.). Notes on thermodynamics. 2d edition. Part I. New York, Wiley, 1901. 12mo. 3 + 69 pp. Cloth. $1.00

SPOONER (H. J.). Elements of geometrical drawing; an elementary text-book on practical plane geometry; including an introduction to solid geometry; written to include the requirements of the syllabus of the Board of education in geometrical drawing for the use of students preparing for the military entrance examinations. New York, Longmans, 1901. 12mo. 39 + 298 pp. Cloth. $1.10

STEINMETZ (C. P.). Theoretical elements of electrical engineering. London, Whittaker, 1901. 8vo. 320 pp. Cloth. 12s. 6d.

STRECKER (K.). See THOMPSON (S. P.).

THOMPSON (S. P.). Die dynamoelektrischen Maschinen. 6te Auflage. Nach C. Grawinkels Uebersetzung neu bearbeitet von K. Strecker und F. Vesper. Halle, Knapp, 1901. 8vo. 11 + 806 pp., 22 plates. M. 24.00

VESPER (F.). See THOMPSON (S. P.).

VUIBERT (H.). Problèmes de baccalauréat (mathématiques), à l'usage des candidats aux baccalauréats de l'enseignement secondaire classique et de l'enseignement secondaire moderne. 3e édition. Paris, Nony, 1901. 8vo. 531 pp.

WILCZYNSKI (E. J.). An application of group theory to hydrodynamics. 1900. 4to. (*Transactions of the American Mathematical Society* 1, pp. 339–352.)

WEISBACH (J.). Lehrbuch der Ingenieur- und Maschinenmechanik. (In 3 Teilen.) Teil 3: Die Mechanik der Zwischen- und Arbeitsmaschinen. 2te, umgearbeitete und vervollständigte Auflage, bearbeitet von G. Herrmann. Abteilung 3: Die Maschinen zur Formveränderung. 2te Hälfte. Braunschweig, Vieweg, 1901. 8vo. Pp. 1–7 and 1223–2200. M. 27.00

ZEUNER (G.). Technische Thermodynamik. 2te Auflage; zugleich 4te Auflage der "Grundzüge der mechanischen Wärmetheorie." Vol. II: Die Lehre von den Dämpfen. Leipzig, Felix, 1901. 8vo. 8 + 463 + 29 pp. M. 14.00

ZIMMERMANN. Ueber Raumfachwerke. Neue Formen und Berechnungsweisen für Kuppeln und sonstige Dachbauten. Berlin, Ernst, 1901. 8vo. 6 + 93 pp. M. 8.00